


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/11, C12Q 1/68, C07H 21/02		A1	(11) International Publication Number: WO 99/33971
			(43) International Publication Date: 8 July 1999 (08.07.99)
(21) International Application Number: PCT/US98/27728 (22) International Filing Date: 30 December 1998 (30.12.98) (30) Priority Data: 60/070,278 31 December 1997 (31.12.97) US (71) Applicant: MEDICAL COLLEGE OF GEORGIA RE- SEARCH INSTITUTE, INC. [US/US]; 1120 15th Street, Augusta, GA 30912-4810 (US). (72) Inventors: DYNAN, William, S.; 428 Wateroak Lane, Au- gusta, GA 30907 (US). YOO, Sunghan; 2902 Shadow Ridge Drive, Augusta, GA 30909 (US). (74) Agents: PABST, Patrea, L. et al.; Arnall Golden & Gregory, LLP, 2800 One Atlantic Center, 1201 West Peachtree Street, Atlanta, GA 30309-3450 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	

(54) Title: OLIGOMERS THAT BIND TO KU PROTEIN



(57) Abstract

Disclosed are oligomers that bind Ku protein. These oligomers are useful for inhibiting activation of DNA-PK, treating certain forms of autoimmune disease, detection and purification of Ku protein, and identification of proteins that interact with Ku protein. Preferably, the oligomers are composed of nucleotides, nucleotide analogs, or a combination. Most preferably, the oligomers are composed of ribonucleotides. Also disclosed is a method of inhibiting DNA repair, a method of identifying cellular proteins that interact with Ku protein, and a method of treating autoimmune disease in patients with anti-Ku antibodies. The disclosed oligomers can have several preferred features, either alone or in combination, in addition to Ku binding. One such feature, referred to herein as inhibition activity, is inhibition of DNA-PK kinase activity. Another preferred feature, referred to herein as aptamer motifs, is the presence of one or more of the base sequences GCUUCCCCANNAC, A(A/C)AUGA, and AACUUCGA.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

OLIGOMERS THAT BIND TO KU PROTEIN

Field of the Invention

The present invention is generally in the area of oligomers, and more specifically in the area of nucleotide sequences that block the function of

5 DNA-dependent protein kinase.

Statement Regarding Federally Funded Research

The U.S. Federal Government has certain rights in this invention by virtue of grant number GM 35866 to William S. Dynan from the U.S. Public Health Service.

10 Background of the Invention

Ku protein, a heterodimer of 70 kDa and 83 kDa polypeptides, is the regulatory component of the DNA-dependent protein kinase (DNA-PK). Ku protein binds to DNA discontinuities and is essential for DNA double-strand break repair.

15 Ku protein was first identified as an autoantigen in sera from certain patients with autoimmune disease (Mimori et al., J. Biol. Chem. 261, 2274-2278 (1986)). Subsequent characterization showed that Ku protein binds avidly to double-stranded DNA ends and other structural discontinuities in DNA such as nicks, gaps, and hairpins (Mimori & Hardin, J. Biol. Chem. 261, 10375-10379 (1986); Paillard & Strauss, Nucleic Acids Res. 19, 5619-5624 (1991); Zhang & Yaneva, Biochem. Biophys. Res. Commun. 186, 574-479 (1992); Blier et al., J. Biol. Chem. 268, 7594-7601 (1993); Falzon et al., J. Biol. Chem. 268, 10546-10552 (1993)). Further biochemical analysis demonstrated that Ku protein is the regulatory component of the DNA-25 dependent protein kinase (Dvir et al., Proc. Natl. Acad. Sci. USA 89, 11920-11924 (1992); Gottlieb & Jackson, Cell 72, 131-142 (1993)). In the presence of DNA ends, Ku protein can interact with the catalytic subunit of DNA-PK (DNA-PKcs) which is thereby targeted to the DNA. The ability of Ku protein to interact with DNA ends suggested that Ku and DNA-PKcs may30 play a role in DNA repair and recombination (Anderson, Trends Biochem. Sci. 18, 433-437 (1993)). Subsequent characterization of ionizing radiation-sensitive mutant cell lines showed that Ku protein and DNA-PKcs are essential for repair of DNA double-strand breaks and for V(D)J

recombination (Getts & Stamato, *J. Biol. Chem.* 269, 15981-15984 (1994); Rathmell, *Proc. Natl. Acad. Sci. USA* 91, 7623-7627 (1994); Smider et al., *Science* 266, 288-291 (1994); Taccioli et al., *Science* 265, 1442-1445 (1994); Blunt et al., *Cell* 80, 813-823 (1995); Boubnov et al., *Proc. Natl. Acad. Sci. USA* 92, 890-894 (1995); Kirchgessner et al., *Science* 267, 1178-1183 (1995); Lees-Miller et al., *Science* 267, 1183-1185 (1995); Peterson et al., *Proc. Natl. Acad. Sci. USA* 92, 3171-3174 (1995)).

The binding of Ku protein to double-stranded DNA ends is largely sequence-independent. The ability of Ku protein to undergo facilitated transfer between DNA fragments with cohesive ends suggests that Ku protein may be able to interact transiently with two DNAs simultaneously, perhaps serving to align the ends for ligation (Bliss & Lane, *J. Biol. Chem.* 272, 5765-5773 (1997)). Consistent with this, recent atomic force microscopy and electron microscopy studies show images of Ku protein tethering DNA fragments together and participating in loop structures (Cary, *Proc. Natl. Acad. Sci. USA* 94, 4267-4272 (1997); Pang et al., *Cancer Res.* 57, 1412-1415 (1997)). There have also been a number of reports of possible sequence-specific binding of Ku protein to DNA (for example, Knuth et al., *J. Biol. Chem.* 265, 17911-17920 (1990); Messier et al., *Proc. Natl. Acad. Sci. USA* 90, 2685-2689 (1993); Okumura et al., *FEBS Lett.* 356, 94-100 (1994); Roberts et al., *Proc. Natl. Acad. Sci., USA* 91, 6354-6358 (1994)). Most recently, a sequence in the long terminal repeat of mouse mammary tumor virus has been characterized that appears to allow interaction of Ku protein with DNA in the absence of ends or single-stranded regions (Giffin et al., *Nature* 380, 265-268 (1996); Giffin et al., *J. Biol. Chem.* 272, 5647-5658 (1997)).

There is some evidence that Ku protein interacts with RNA, although this has been much less studied than the interaction with DNA. Antibodies to Ku protein stain both the nucleoplasm and the nucleolus. The amount of Ku protein in the nucleolus changes depending on the growth state of the cell, suggesting that this localization is actively regulated (Yaneva & Jhiang, *Biochim. Biophys. Acta* 1090, 181-187 (1991)). Separately, it has been demonstrated that nucleolar staining is sensitive to RNase treatment, whereas

nucleoplasmic staining is not (Reeves, J. Exp. Med. 161, 18-39 (1985)). Thus, nucleolar localization may be regulated by interaction of Ku protein with RNA. Ku protein does not appear to bind to bulk tRNA or to synthetic RNA polymers (Mimori & Hardin, J. Biol. Chem. 261, 10375-10379 (1986)). However, one study showed that Ku protein forms a specific complex with an RNA that included the HIV trans-activation response (TAR) element sequence (Kaczmariski & Khan, Biochem. Biophys. Res. Commun. 196, 935-942 (1993)).

The Ku protein has been suggested to be involved in many important nuclear processes, including transcription, replication, and growth control, as well as DNA repair. It would be useful to have a means to demonstrate whether Ku protein is required for a biochemical activity (such as repair or recombination) that has been reconstituted in a crude cell-free system. It would also be useful to have a means to identify nuclear proteins that physically interact with Ku protein to exert their biological functions.

Ku protein and DNA-PK are important in the repair of radiation-induced DNA damage. If damage cannot be repaired, cells die. The cytotoxic effect of ionizing radiation forms the basis for radiation therapy, which is widely used in the treatment of human cancer. The efficacy of radiation therapy is currently limited by the radiation resistance of certain tumors (for example, glioblastomas) and by the side effects caused by irradiation of nearby normal tissues (for example, in treatment of breast and cervical cancer). Therefore, it would also be useful to have a means for sensitizing target cells and tissues to therapeutic radiation.

Some patients with autoimmune diseases such as systemic lupus erythematosus (SLE) and scleroderma produce anti-Ku antibodies. Ku is one of a number of proteins that are targets of autoantibodies in these patients. High levels of autoantibodies lead to deleterious consequences for the patient. It would be useful to have a compound directed against Ku protein to alter the course of autoimmune disease in patients with anti-Ku antibodies.

A problem in gene therapy is a loss of the foreign DNA from illegitimate recombination. It is believed that illegitimate recombination

requires Ku-dependent double strand break repair. It would be useful to have a means to improve the stability of transgene DNA.

Therefore, it is an object of the disclosed invention to provide oligomers that bind to Ku protein.

5 It is also an object of the disclosed invention to provide oligomers that prevent illegitimate recombination to improve the stability of transgene DNA.

It is another object of the disclosed invention to provide oligomers that recognize Ku protein in a complex environment containing other
10 macromolecules.

It is also an object of the disclosed invention to provide oligomers that improve the efficacy of radiation therapy by inhibiting DNA repair in the target cells.

It is also an object of the disclosed invention to provide oligomers to
15 treat autoimmune disease in patients with anti-Ku antibodies.

It is also an object of the disclosed invention is to provide an assay for cellular proteins that interact with Ku protein or that promote or inhibit the interaction between Ku protein and DNA-PKcs.

It is also an object of the disclosed invention is to provide oligomers
20 useful for manipulating the activity of Ku protein in cells and organisms to better understand its physiological role.

Summary of the Invention

Disclosed are oligomers that bind Ku protein. These oligomers, also referred to herein as aptamers, are useful for inhibiting activation of DNA-
25 PK, treating certain forms of autoimmune disease, detection and purification of Ku protein, and identification of proteins that interact with Ku protein. Preferably, the oligomers are composed of nucleotides, nucleotide analogs, or a combination. Most preferably, the oligomers are composed of ribonucleotides. Also disclosed is a method of inhibiting DNA repair, a
30 method of identifying cellular proteins that interact with Ku protein, and a method of treating autoimmune disease in patients with anti-Ku antibodies.

The disclosed oligomers can have several preferred features, either alone or in combination, in addition to Ku binding. One such feature,

referred to herein as inhibition activity, is inhibition of DNA-PK kinase activity. As discussed above, interaction of the Ku protein and DNA is involved in activation of DNA-PK kinase activity. Another preferred feature, referred to herein as aptamer motifs, is the presence of one or more

5 of the base sequences GCUUUCCCCANNAC, A(A/C)AUGA, and AACUUCGA. These sequences -- referred to herein as aptamer motif 1, aptamer motif 2, and aptamer motif 3, respectively -- are associated with Ku binding capability. Another preferred feature, referred to herein as aptamer structure, is the presence of a structure similar to the structure shown in

10 Figure 6A. This structure has the general formula 5'-A-B-C-D-C'-E-A'-3', where A, B, C, D, C', E, and A' are components of the oligomer. In this structure, A and A' interact to form a stem structure, C and C' interact to form a stem structure, B and E make up a bulge region, and D is either a bulge or a loop. Figure 6A depicts component D as a loop. Each of these

15 preferred features (inhibition activity, aptamer motif, and aptamer structure) can be used either alone or in combination with one or both of the other characteristics.

Brief Description of the Drawings

Figure 1 is a graph of percent RNA or DNA bound versus

20 concentration of Ku (nM) for five representative Ku protein-RNA and DNA binding curves obtained with an electrophoretic mobility shift assay. The graph shows that pooled RNA tested after the fourth and sixth round of Systematic Evolution of Ligands by Exponential Enrichment (SELEX; Tuerk & Gold, Science 249:505-510 (1990), and U.S. Patent Numbers 5,270,163

25 and 5,567,588) had an increased ability to bind to Ku protein. Radiolabeled nucleic acid probes were as follows: □, nonselected RNA; ▽, RNA after 4th round of selection; ▲, RNA after 6th round of selection; O, HIV-TAR RNA (included for comparison); ◆, 21 base pair double-stranded DNA.

Figure 2A is a graph of kinase activity (percentage of activity in

30 absence of RNA) versus RNA concentration (nM) for eight representative aptamers. The aptamers were as follows: X, nonselected RNA; O, #2; □, #7-3; △, SC4; +, SC5; ◆, SC8; ▽, #1-2; ●, SC2.

Figure 2B is a graph of DNA binding (percent of Ku-DNA complexes detected in the absence of competitor) versus RNA concentration (nM) for seven representative aptamers. Purified Ku protein (2 nM) was incubated with radiolabeled 21 base pair double-stranded DNA (1 nM) in the presence of various amounts of nonselected RNA or aptamer. The aptamers were as follows: X, nonselected RNA; O, #2; □, #7-3; △, SC4; +, SC5; ◆, SC8; ▼, #1-2; ●, SC2.

Figure 3 is a bar graph of kinase activity (percentage of activity in the absence of RNA) in the presence of different aptamers, each at three different concentrations. Reactions were performed in crude nuclear extract. Reactions contained 2 µl HeLa cell nuclear extract (0.4 µg protein), substrate (Ku protein (2 nM)) and DNA (1 nM) were as in Figure 2. Reactions were performed in the presence of various amounts of aptamer (250, 500, 1000 nM).

Figure 4 is the predicted secondary structure of examples of the disclosed aptamers, SC4 and #2. RNA secondary structures were predicted using the methods of Zuker and co-workers (Jaeger et al., 1989; Jaeger et al., 1990) as implemented on the mfold World Wide Web server (www.ibc.wustl.edu/zuker/rna). Boldface indicates the conserved sequence motifs as shown in Table 1.

Figure 5 shows the predicted secondary structure of the aptamer SC4 and the predicted structures of the mutated aptamers, SC4/D1 and SC4/D2. The 3' invariant region of the aptamer, i.e. the twenty-three 3' terminal bases, was removed from the SC4/D1 aptamer and an additional 15 bases were removed from the 3' end of the molecule in the SC4/D2 aptamer.

Figure 6A is a representation of the general secondary structure for a preferred form of aptamer.

Figure 6B is a comparison of the predicted secondary structure of a preferred form of aptamer with the secondary structure of a hypothetical double-strand DNA break repair intermediate.

Detailed Description of the Invention

Disclosed are oligomers that bind Ku protein. It is preferred that the oligomers bind Ku protein with a dissociation constant of less than 10 nM.

These oligomers are useful for inhibiting activation of DNA-PK, treating certain forms of autoimmune disease, detection and purification of Ku protein, and identification of proteins that interact with Ku protein. Preferably, the oligomers are composed of nucleotides, nucleotide analogs, or a combination. Most preferably, the oligomers are composed of ribonucleotides. Also disclosed is a method of inhibiting DNA repair, a method of identifying cellular proteins that interact with Ku protein, and a method of treating autoimmune disease in patients with anti-Ku antibodies.

The disclosed oligomers can have several preferred features, either alone or in combination, in addition to Ku binding. One such feature, referred to herein as inhibition activity, is inhibition of DNA-PK kinase activity. As discussed above, interaction of the Ku protein and DNA is involved in activation of DNA-PK kinase activity. Another preferred feature, referred to herein as aptamer motifs, is the presence of one or more of the base sequences GCUUCCCCANNAC, A(A/C)AUGA, and AACUUCGA. These sequences -- referred to herein as aptamer motif 1, aptamer motif 2, and aptamer motif 3, respectively -- are associated with Ku binding capability. Another preferred feature, referred to herein as aptamer structure, is the presence of a structure similar to the structure shown in Figure 6A. This structure has the general formula 5'-A-B-C-D-C'-E-A'-3', where A, B, C, D, C', E, and A' are components of the oligomer. In this structure, A and A' interact to form a stem structure, C and C' interact to form a stem structure, B and E make up a bulge region, and D is either a bulge or a loop. Figure 6A depicts component D as a loop. Each of these preferred features (inhibition activity, aptamer motif, and aptamer structure) can be used either alone or in combination with one or both of the other characteristics.

Examples of preferred oligomers are listed in Table 1. These oligoribonucleotides were identified as described in the examples. Only the portion of the selected oligomers that arose from the variable region of the pool RNA is shown in Table 1. That is, each of the oligonucleotides listed in Table 1 include the sequence GGGAGGAUAUUUCUCAGACCGUAA at the 5' end and the sequence UUGCAGCAUCGUGAACUAGGAUC at the

3' end, although these sequences are not shown in Table 1. Thus, oligomer SC6 (SEQ ID NO:1) has the sequence GGGAGGAUAUUUUCUCAGACCGUAAGACUCACGAUGGACCAUACGCCUUCCCACUGGUCUUGUUAUUGCAGCAUCGUGAACUAGGAUC, although only the sequence
5 GACUCACGAUGGACCAUACGCCUUCC CACUGGUCUUGUUA is shown in the table.

Table 1 also discloses the dissociation constants (Kd) of the individual aptamers and the ability of the individual aptamers to inhibit DNA-PK activity. In a standard reaction, aptamers with a DNA-PK activity
10 denoted as A had more than 85% inhibition, aptamers denoted as B had between 30 and 80% inhibition, and aptamers denoted with a C had under 30% inhibition.

A. Oligomers

Some embodiments of the oligomers can be described generally by
15 an oligonucleotide comprising the components A, B, C, D, C', E, and A'. Component A is the 5' most component and component A' is the 3' most component of the Ku binding oligonucleotides. Components A and A' interact in such a way as to form a stem structure through nucleotide base pairing. It is not required that all of the base pairs be contiguous, but it is
20 recognized that those of skill in the art understand that the stability of any given double stranded nucleic acid molecule is a function of both the number of base pairs present and the number of bulges or mismatches that are present. A preferred embodiment has a stem that has a level of stability equivalent to two C:G base pairs. Another preferred embodiment has a
25 purine:purine base pair, three base pairs from the bulge region comprised by components B and E (see Figure 4).

Component B represents a bulge structure contiguous and 3' to component A. Furthermore, this component is preferably composed of pyrimidines. Most preferably this component contains the sequence 5'-
30 UUUCCC-3' or 5'-UUUCCCA-3'.

Table 1: RNA Sequences and K_d values from the SELEX Procedure

Name Frequency	Aligned sequence	DNA-PK Inhibition	K_d
CLASS I			
SC6 (3)	GACUACGAUGGACCAUAGCGCUUCCCAUGGUCUUGUUA (SEQ ID NO:1)	C	2.0 nM
#1-2 (1)	CAACACCUUGCUUCCCAUACCCUGAAUACAGUCGGAU (SEQ ID NO:2)	A	1.5 nM
#1-17 (1)	UCCUUAUUUUAUGGCUUCCCAAGCAGACAAAGCUCUCGG (SEQ ID NO:3)	B	3.2 nM
#85 (1)	CAAGUAUCACGCAUUCUCCCAUUCACUGUAGAGACUGA (SEQ ID NO:4)	B	0.7 nM
HYBRID			
#2 (1)	GCCUAUGCACGGAGCUUCCCGAGCUACAGAUAGAAACCAGC (SEQ ID NO:5)	A	0.3 nM
SC4 (8)	CCUAGUCUAAUCCGAGGCUUCCCGAGGAC_AAUAGACCCAC (SEQ ID NO:6)	A	1.7 nM
SC5 (4)	CUUGAACAUAGAUAG_GCUUACCCAUAGACAGAUUGACCCUU (SEQ ID NO:7)	A	2.0 nM
CLASS II			
SC9 (3)	UGCCUUUAGCUGCGAC_AAUGAACAGCAUGACCUCACUAC (SEQ ID NO:8)	B	0.6 nM
SC8 (8)	(SEQ ID NO:9) GUCCUUACAUAAUGCUUACAGACACACUAAGAACGUCAC	A	0.6 nM
SC3 (3)	CAUACACACAGUUCUAGCAUCCCGCAAUGGUAAGUCCGCA (SEQ ID NO:10)	B	0.8 nM
SC1 (23)	UUGUUCACCCUUGUCUAAACAUGAUACCGAUACGGACUACA (SEQ ID NO:11)	B	1.2 nM
#84 (1)	AUCCGCGUACCGGCUCAAUUGUCACUUAUAGUAGAAAGCA (SEQ ID NO:12)	B	1.8 nM
#52 (1)	(SEQ ID NO:13) CUGAUCGUUCAAUAGAUAUUCUUUACCUUGAGUAAACCGA	C	3.2 nM
HYBRID			
SC12 (4)	(SEQ ID NO:14) CUCGCAACAUGACUUCGAAAGUUUAAUUCGUUCUUGUCAA	B	0.5 nM
CLASS III			
#7-3 (1)	AGGUCGGCAUACAGAGUUCGCCGAUUGCGCGUAAACUUCGACU (SEQ ID NO:15)	B	1.8 nM
SC11 (2)	(SEQ ID NO:16) CUUAGUUUCGUAUCGAAGCUCAUUGGCCACGGUGGAUAAC	A	>10 nM
SC2 (12)	CACGCUCUACAAACAGAUUGCGGAUUAAACUUCGCUUCAUA (SEQ ID NO:17)	A	0.8 nM
#42 (1)	(SEQ ID NO:18) CAUCCUGGUACUCACUUCGACAUUCGUACGUUCAUUAUAC	B	4.5 nM
OTHER			
SC13 (3)	ACCUUUUUAGACGAACCUCAAGAAAGUACAUAUUAUUGUAGAAAC (SEQ ID NO:19)	B	0.8 nM

RNA sequences are aligned to show maximum sequence identity. The underlined sequence indicates sequence identity within class.

Component C is contiguous and 3' to component B. Components C and C' form a stem region between the bulge formed by components B and E and the loop region D. It is also understood that this region can contain mismatches and bulges but that a certain level of stability is required. In preferred embodiments this stem will possess a stability equivalent to a stem made up of an AGCU:UCGA stem.

Component D is contiguous and 3' to component C. Component C' is contiguous and 3' to component D. Component D is a sequence which preferably forms a loop structure. In a preferred embodiment component D contains the sequence 5'-ACNNAUGA-3'. In another preferred embodiment component D contains the sequence 5'-ACNNAUGANNNN-3'. In another preferred embodiment component D contains 5'-ACMAUGANNNN-3' (where M is A or C). In a most preferred embodiment Component D contains the sequence 5'-ACAGAUGAAACC-3'. It is also recognized that this loop structure can also be represented by a bulge-stem structure. Those in the art will recognize how to turn the loop structure into a bulge-stem structure. This would be done for example if there were two oligomers that were used to form the preferred structure.

Component E is contiguous and 3' to component C'. Component E is a bulged region. In the most preferred embodiment component B contains the sequence 5'-UGC-3'.

1. Oligomers Containing Chemically Modified Nucleotides.


RNA is relatively labile and can be degraded by a number of ribonucleases. This degradation can be greatly reduced by the introduction of modifications and substitutions at the 2'-prime position of the ribonucleotide and by modifications and substitutions along the phosphate backbone of the RNA. In addition a variety of modifications can be made on the nucleobases themselves which both inhibit degradation and which can increase desired nucleotide interactions or decrease undesired nucleotide interactions. Once the sequence of an aptamer is known modifications or substitutions can be made by the synthetic procedures described below or by

procedures known to those of skill in the art. Aptamers with these various modifications can then be tested for function using any suitable assay for the Ku function of interest, such as the Ku protein binding assay or the inhibition of DNA-PK activity assay.

5 The oligomers disclosed are made up of nucleotides and/or nucleotide analogs or a combination of both, or are oligonucleotide analogues. The disclosed oligomers described by the general structure above may contain nucleotide analogs at positions which do not effect the function of the oligomer to bind Ku protein. The oligomeric sequences specifically interact
10 with Ku protein, preferably at the DNA binding site of Ku protein.

Components A, A', B, C, C', D, and E are made up of nucleotides and nucleotide analogues in which the nucleotide and nucleotide analogs have the structure

15



(I)

20

25 In structure (I) each S can be adenin-9-yl, cytosin-1-yl, guanin-9-yl, uracil-1-yl, uracil-5-yl, hypoxanthin-9-yl, thymine-1-yl, 5-methylcytosin-1-yl, 2,6-diaminopurin-9-yl, purin-9-yl, 7-deazaadenin-9-yl, 7-deazaguanin-9-yl, 5-propynylcytosin-1-yl, 5-propynyluracil-1-yl, isoguanin-9-yl, 2-aminopurin-9-yl, 6-methyluracil-1-yl, 4-thiouracil-1-yl, 2-pyrimidone-1-yl,
30 quinazoline-2,4-dione-1-yl, xanthin-9-yl, N²-dimethylguanin-9-yl or a functional equivalent thereof;

Each V can be an O, S, NH, or CH₂ group.

Each W can be -H, -OH, -COOH, -CONH₂, -CONHR¹, -CONR¹R², -NH₂, -NHR¹, -NR¹R², -NHCOR¹, -SH, SR¹, -F, -ONH₂, -ONHR¹, -ONR¹R², -NHOH, -NHOR¹, -NR²OH, -NR²OR¹, substituted or unsubstituted C₁-C₁₀ straight chain or branched alkyl, substituted or
 5 unsubstituted C₂-C₁₀ straight chain or branched alkenyl, substituted or unsubstituted C₂-C₁₀ straight chain or branched alkynyl, substituted or unsubstituted C₁-C₁₀ straight chain or branched alkoxy, substituted or unsubstituted C₂-C₁₀ straight chain or branched alkenyloxy, and substituted or unsubstituted C₂-C₁₀ straight chain or branched alkynyloxy. The
 10 substituents for W groups are independently halogen, cyano, amino, carboxy, ester, ether, carboxamide, hydroxy, or mercapto. R¹ and R² can be substituted or unsubstituted alkyl, alkenyl, or alkynyl groups, where the substituents are independently halogen, cyano, amino, carboxy, ester, ether, carboxamide, hydroxy, or mercapto.

15 T and Z are residues which together form a phosphodiester or phosphorothioate diester bond between adjacent nucleosides or nucleoside analogues or together form an analogue of an internucleosidic bond.

The 3' end of the disclosed oligomers can be protected against degradation by exonucleases by, for example, using a nucleotide analogue
 20 that is modified at the 3' position of the ribose sugar (for example, by including a substituted or unsubstituted alkyl, alkoxy, alkenyl, alkenyloxy, alkynyl or alkynyloxy group as defined above). The disclosed oligomers can also be stabilized against degradation at the 3' end by exonucleases by including a 3'-3'-linked dinucleotide structure (Ortigao *et al.*, *Antisense Research and Development* 2:129-146 (1992)) and/or two modified phospho
 25 bonds, such as two phosphorothioate bonds.

The disclosed oligomers can also be linked to a prosthetic group in order to improve their cellular uptake and/or to enable a specific cellular localization. Examples of such prosthetic groups are polyamino acids (for
 30 example, polylysine), lipids, hormones or peptides. These prosthetic groups are usually linked via the 3' or 5' end of the oligomer either directly or by means of suitable linkers (for example, linkers based on 6-aminohexanol or

6-mercaptohexanol). These linkers are commercially available and techniques suitable for linking prosthetic groups to the oligomer are known to a person skilled in the art. The oligomers can also be produced as fusion oligonucleotides as discussed below.

5 In preferred oligomers, the oligonucleotides are made from RNA. The examples of Ku aptamers obtained as described in the examples are in the form of RNA. These RNAs were selected from a pool of about 10^{14} different RNA sequences using the systematic evolution of ligands by exponential enrichment (SELEX) procedure. Most of the selected RNAs
10 bind to Ku protein with an equilibrium dissociation constant (K_d) ≤ 2 nM, comparable to the affinity of DNA fragments for Ku protein under similar conditions. Many of the RNAs inhibited DNA-PK activity up to 85% by competing with DNA for a common binding site in Ku protein. None of the several RNAs tested activated DNA-PK in the absence of DNA.

15 2. Isolating RNAs That Bind Ku Protein.

 There are several techniques that can be adapted for refinement of the disclosed aptamers or the selection of additional aptamers. Preferred are the techniques described in the examples. Other techniques can also be used. These include techniques generally referred to as "in vitro genetics"
20 (Szostak, TIBS 19:89, 1992).

 Molecules that inhibit or activate Ku mediated DNA-PK function can be isolated by screening for molecules that inhibit function of DNA-PK without inhibiting the interaction of DNA-PK with Ku protein. For example, molecules that inhibit DNA binding to DNA-PK may inhibit DNA-PK
25 function without inhibiting DNA-PK:Ku interaction.

 The disclosed aptamers can also be refined through the use of computer modeling techniques. Examples of molecular modeling systems are the CHARMM and QUANTA programs, Polygen Corporation, Waltham, MA. CHARMM performs the energy minimization and molecular dynamics
30 functions. QUANTA performs the construction, graphic modeling and analysis of molecular structure. QUANTA allows interactive construction, modification, visualization, and analysis of the behavior of molecules with

each other. These applications can be adapted to define and display the secondary structure of RNA and DNA molecules.

The pool of nucleic acid molecules from which the disclosed aptamers were isolated included invariant sequences flanking a variable
5 sequence of approximately forty nucleotides. The 5' invariant region contained 27 nucleotides and the 3' invariant region contained 25 nucleotides. The invariant regions were used for the amplification of the molecules that were enriched through the SELEX procedure described below. This pool was chosen because it had been shown to function well in
10 SELEX protocols. (Chen and Gold, *Biochemistry* 33, 8746-8756 (1994)).

3. Synthesis of Oligomers.

The disclosed oligomers can be synthesized using any suitable method. Many synthesis methods are known. The following techniques are preferred for synthesis of for example, 2'-O-Allyl modified oligomers that
15 contain residual purine ribonucleotides, and bearing a suitable 3'-terminus such as an inverted thymidine residue (Ortigao et al., *Antisense Research and Development* 2:129-146 (1992)) or two phosphorothioate linkages at the 3'-terminus to prevent eventual degradation by 3'-exonucleases, can be synthesized by solid phase β -cyanoethyl phosphoramidite chemistry (Sinha
20 et al., *Nucleic Acids Res.* 12:4539-4557 (1984)) on any commercially available DNA/RNA synthesizer. A preferred method is the 2'-O-*tert*-butyldimethylsilyl (TBDMS) protection strategy for the ribonucleotides (Usman et al., *J. Am. Chem. Soc.* 109:7845-7854 (1987)), and all the required 3'-O-phosphoramidites are commercially available. In addition, the
25 use of aminomethylpolystyrene is preferred as the support material due to its advantageous properties (McCollum and Andrus *Tetrahedron Letters* 32:4069-4072 (1991)). Fluorescein can be added to the 5'-end of a substrate RNA during the synthesis by using commercially available fluorescein phosphoramidites. In general, a desired oligomer can be synthesized using a
30 standard RNA cycle. Upon completion of the assembly, all base labile protecting groups are removed by an 8 hour treatment at 55°C with concentrated aqueous ammonia/ethanol (3:1 v/v) in a sealed vial. The

ethanol suppresses premature removal of the 2'-O-TBDMS groups which would otherwise lead to appreciable strand cleavage at the resulting ribonucleotide positions under the basic conditions of the deprotection (Usman *et al.*, *J. Am. Chem. Soc.* 109:7845-7854 (1987)). After

5 lyophilization the TBDMS protected oligomer is treated with a mixture of triethylamine trihydrofluoride/triethylamine/N-methylpyrrolidinone for 2 hours at 60°C to afford fast and efficient removal of the silyl protecting groups under neutral conditions (Wincott *et al.*, *Nucleic Acids Res.* 23:2677-2684 (1995)). The fully deprotected oligomer can then be precipitated with

10 butanol according to the procedure of Cathala and Brunel (*Nucleic Acids Res.* 18:201 (1990)). Purification can be performed either by denaturing polyacrylamide gel electrophoresis or by a combination of ion-exchange HPLC (Sproat *et al.*, *Nucleosides and Nucleotides* 14:255-273 (1995)) and reversed phase HPLC. For use in cells, it is preferred that synthesized

15 oligomers be converted to their sodium salts by precipitation with sodium perchlorate in acetone. Traces of residual salts are then preferably removed using small disposable gel filtration columns that are commercially available. As a final step it is preferred that the authenticity of the isolated oligomers is checked by matrix assisted laser desorption mass spectrometry (Pieles *et al.*,

20 *Nucleic Acids Res.* 21:3191-3196 (1993)) and by nucleoside base composition analysis. In addition, a functional cleavage test with the oligomer on the corresponding chemically synthesized short oligoribonucleotide substrate is also preferred.

The disclosed oligomers can also be produced through enzymatic

25 methods, when the nucleotide subunits are available for enzymatic manipulation. For example, the RNA molecules can be made through *in vitro* RNA polymerase T7 reactions. They can also be made by strains of bacteria or cell lines expressing T7, and then subsequently isolated from these cells. As discussed below, the disclosed aptamers can also be

30 expressed in cells directly using vectors and promoters.

B. Pharmaceutical Compositions.

The disclosed oligomers can be used in pharmaceutical mixtures that contain one or several oligomers as the active substance, and, optionally, pharmaceutically acceptable auxiliary substances, additives and carriers.

- 5 Such pharmaceutical oligomers are suitable for the production of an agent to specifically inhibit Ku DNA binding activity in cells or inhibit the activity of DNA-PK. The disclosed oligomers can also be used for Ku inhibition and DNA-PK inhibition in plants or animals. Thus, the disclosed oligomers are expected to be useful as drugs for humans and animals as well as a pesticide
10 for plants.

As used herein, microparticles include liposomes, virosomes, microspheres and microcapsules formed of synthetic and/or natural polymers. Methods for making microcapsules and microspheres are known to those skilled in the art and include solvent evaporation, solvent casting, spray drying and solvent
15 extension. Examples of useful polymers which can be incorporated into various microparticles include polysaccharides, polyanhydrides, polyorthoesters, polyhydroxides and proteins and peptides.

Liposomes can be produced by standard methods such as those reported by Kim *et al.*, *Biochim. Biophys. Acta*, 728:339-348 (1983); Liu *et al.*, *Biochim. Biophys. Acta*, 1104:95-101 (1992); and Lee *et al.*, *Biochim. Biophys. Acta*, 1103:185-197 (1992); Wang *et al.*, *Biochem.*, 28:9508-9514 (1989)). Such methods have been used to deliver nucleic acid molecules to the nucleus and cytoplasm of cells of the MOLT-3 leukemia cell line (Thierry and Dritschilo, *Nucl. Acids Res.*, 20:5691-5698 (1992)).
20

- 25 Alternatively, the disclosed oligomers can be incorporated within microparticles, or bound to the outside of the microparticles, either ionically or covalently.

Cationic liposomes or microcapsules are microparticles that are particularly useful for delivering negatively charged compounds such as the disclosed compounds, which can bind ionically to the positively charged
30 outer surface of these liposomes. Various cationic liposomes have previously been shown to be very effective at delivering nucleic acids or

nucleic acid-protein complexes to cells both *in vitro* and *in vivo*, as reported by Felgner *et al.*, *Proc. Natl. Acad. Sci. USA*, 84:7413-7417 (1987); Felgner, *Advanced Drug Delivery Reviews*, 5:163-187 (1990); Clarenc *et al.*, *Anti-Cancer Drug Design*, 8:81-94 (1993). Cationic liposomes or microcapsules
5 can be prepared using mixtures including one or more lipids containing a cationic side group in a sufficient quantity such that the liposomes or microcapsules formed from the mixture possess a net positive charge which will ionically bind negatively charged compounds. Examples of positively charged lipids that may be used to produce cationic liposomes include the
10 aminolipid dioleoyl phosphatidyl ethanolamine (PE), which possesses a positively charged primary amino head group; phosphatidylcholine (PC), which possess positively charged head groups that are not primary amines; and N[1-(2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium ("DOTMA," see Felgner *et al.*, *Proc. Natl. Acad. Sci. USA*, 84:7413-7417 (1987); Felgner *et al.*, *Nature*, 337:387-388 (1989); Felgner, *Advanced Drug Delivery Reviews*,
15 5:163-187 (1990)).

The disclosed oligomers can also be encapsulated by or coated on cationic liposomes which can be injected intravenously into a mammal. This system has been used to introduce DNA into the cells of
20 multiple tissues of adult mice, including endothelium and bone marrow, where hematopoietic cells reside (see, for example, Zhu *et al.*, *Science*, 261:209-211 (1993)).

C. Vectors And Expression Sequences

Delivery and expression of the disclosed oligomers can be facilitated
25 by vectors and expression sequences. As used herein, plasmid or viral vectors are agents that transport the gene into the cell without degradation and include a promoter yielding expression of the gene in the cells into which it is delivered. In a preferred embodiment vectors are derived from either a virus or a retrovirus. Preferred viral vectors are Adenovirus, Adeno-associated virus, Herpes virus, Vaccinia virus, Polio virus, AIDS virus,
30 neuronal trophic virus, Sindbis and other RNA viruses, including these viruses with the HIV backbone. Also preferred are any viral families which

share the properties of these viruses which make them suitable for use as vectors. Preferred retroviruses include Murine Maloney Leukemia virus, MMLV, and retroviruses that express the desirable properties of MMLV as a vector. Retroviral vectors are able to carry a larger genetic payload, i.e., a
5 transgene or marker gene, than other viral vectors, and for this reason are a commonly used vector. However, they are not useful in non-proliferating cells. Adenovirus vectors are relatively stable and easy to work with, have high titers, and can be delivered in aerosol formulation, and can transfect non-dividing cells. Pox viral vectors are large and have several sites for
10 inserting genes, they are thermostable and can be stored at room temperature. A preferred embodiment is a viral vector which has been engineered so as to suppress the immune response of the host organism, elicited by the viral antigens. Preferred vectors of this type will carry coding regions for Interleukin 8 or 10.

15 Viral vectors have higher transaction (ability to introduce genes) abilities than do most chemical or physical methods to introduce genes into cells. Typically, viral vectors contain, nonstructural early genes, structural late genes, an RNA polymerase III transcript, inverted terminal repeats necessary for replication and encapsidation, and promoters to control the
20 transcription and replication of the viral genome. When engineered as vectors, viruses typically have one or more of the early genes removed and a gene or gene/promotor cassette is inserted into the viral genome in place of the removed viral DNA. Constructs of this type can carry up to about 8 kb of foreign genetic material. The necessary functions of the removed early
25 genes are typically supplied by cell lines which have been engineered to express the gene products of the early genes in trans.

1. Retroviral Vectors.

A retrovirus is an animal virus belonging to the virus family of Retroviridae, including any types, subfamilies, genus, or tropisms.
30 Retroviral vectors, in general, are described by Verma, I.M., Retroviral vectors for gene transfer. In MICROBIOLOGY-1985, American Society for Microbiology, pp. 229-232, Washington, (1985). Examples of methods for

using retroviral vectors for gene therapy are described in U.S. Patent Nos. 4,868,116 and 4,980,286; PCT applications WO 90/02806 and WO 89/07136; and Mulligan, (Science 260:926-932 (1993)).

5 A retrovirus is essentially a package which has packed into it nucleic acid cargo. The nucleic acid cargo carries with it a packaging signal, which ensures that the replicated daughter molecules will be efficiently packaged within the package coat. In addition to the package signal, there are a number of molecules which are needed in cis, for the replication, and packaging of the replicated virus. Typically a retroviral genome, contains
10 the gag, pol, and env genes which are involved in the making of the protein coat. It is the gag, pol, and env genes which are typically replaced by the foreign DNA that it is to be transferred to the target cell. Retrovirus vectors typically contain a packaging signal for incorporation into the package coat, a sequence which signals the start of the gag transcription unit, elements
15 necessary for reverse transcription, including a primer binding site to bind the tRNA primer of reverse transcription, terminal repeat sequences that guide the switch of RNA strands during DNA synthesis, a purine rich sequence 5' to the 3' LTR that serve as the priming site for the synthesis of the second strand of DNA synthesis, and specific sequences near the ends of
20 the LTRs that enable the insertion of the DNA state of the retrovirus to insert into the host genome. The removal of the gag, pol, and env genes allows for about 8 kb of foreign sequence to be inserted into the viral genome, become reverse transcribed, and upon replication be packaged into a new retroviral particle. This amount of nucleic acid is sufficient for the delivery of a one to
25 many genes depending on the size of each transcript. It is preferable to include either positive or negative selectable markers along with other genes in the insert.

Since the replication machinery and packaging proteins in most retroviral vectors have been removed (gag, pol, and env), the vectors are
30 typically generated by placing them into a packaging cell line. A packaging cell line is a cell line which has been transfected or transformed with a retrovirus that contains the replication and packaging machinery, but lacks

any packaging signal. When the vector carrying the DNA of choice is transfected into these cell lines, the vector containing the gene of interest is replicated and packaged into new retroviral particles, by the machinery provided in cis by the helper cell. The genomes for the machinery are not packaged because they lack the necessary signals.

2. Adenoviral Vectors.

The construction of replication-defective adenoviruses has been described (Berkner et al., J. Virology 61:1213-1220 (1987); Massie et al., Mol. Cell. Biol. 6:2872-2883 (1986); Haj-Ahmad et al., J. Virology 57:267-274 (1986); Davidson et al., J. Virology 61:1226-1239 (1987); Zhang "Generation and identification of recombinant adenovirus by liposome-mediated transfection and PCR analysis" BioTechniques 15:868-872 (1993)). The benefit of the use of these viruses as vectors is that they are limited in the extent to which they can spread to other cell types, since they can replicate within an initial infected cell, but are unable to form new infectious viral particles. Recombinant adenoviruses have been shown to achieve high efficiency gene transfer after direct, *in vivo* delivery to airway epithelium, hepatocytes, vascular endothelium, CNS parenchyma and a number of other tissue sites (Morsy, J. Clin. Invest. 92:1580-1586 (1993); Kirshenbaum, J. Clin. Invest. 92:381-387 (1993); Roessler, J. Clin. Invest. 92:1085-1092 (1993); Moullier, Nature Genetics 4:154-159 (1993); La Salle, Science 259:988-990 (1993); Gomez-Foix, J. Biol. Chem. 267:25129-25134 (1992); Rich, Human Gene Therapy 4:461-476 (1993); Zabner, Nature Genetics 6:75-83 (1994); Guzman, Circulation Research 73:1201-1207 (1993); Bout, Human Gene Therapy 5:3-10 (1994); Zabner, Cell 75:207-216 (1993); Caillaud, Eur. J. Neuroscience 5:1287-1291 (1993); and Ragot, J. Gen. Virology 74:501-507 (1993)). Recombinant adenoviruses achieve gene transduction by binding to specific cell surface receptors, after which the virus is internalized by receptor-mediated endocytosis, in the same manner as wild type or replication-defective adenovirus (Chardonnet and Dales, Virology 40:462-477 (1970); Brown and Burlingham, J. Virology 12:386-396 (1973); Svensson and Persson, J.

Virology 55:442-449 (1985); Seth, et al., J. Virol. 51:650-655 (1984); Seth, et al., Mol. Cell. Biol. 4:1528-1533 (1984); Varga et al., J. Virology 65:6061-6070 (1991); Wickham et al., Cell 73:309-319 (1993)).

A preferred viral vector is one based on an adenovirus which has had
5 the E1 gene removed and these virions are generated in a cell line such as the human 293 cell line. In another preferred embodiment both the E1 and E3 genes are removed from the adenovirus genome.

Another type of viral vector is based on an adeno-associated virus (AAV). This defective parvovirus is a preferred vector because it can infect
10 many cell types and is nonpathogenic to humans. AAV type vectors can transport about 4 to 5 kb and wild type AAV is known to stably insert into chromosome 19. Vectors which contain this site specific integration property are preferred. An especially preferred embodiment of this type of vector is the P4.1 C vector produced by Avigen, San Francisco, CA, which
15 can contain the herpes simplex virus thymidine kinase gene, HSV-tk, and/or a marker gene, such as the gene encoding the green fluorescent protein, GFP.

The inserted genes in viral and retroviral usually contain promoters, and/or enhancers to help control the expression of the desired gene product. A promoter is generally a sequence or sequences of DNA that function when
20 in a relatively fixed location in regard to the transcription start site. A promoter contains core elements required for basic interaction of RNA polymerase and transcription factors, and may contain upstream elements and response elements.

3. Expression Sequences.

25 Expression of the disclosed aptamers in cells also requires sequences to direct expression. It is preferred that an RNA polymerase III (pol III) promoter be used for expression. Pol III promoters generate transcripts that can be engineered to remain in the nucleus of the cell where Ku protein acts. It is preferred that a complete pol III transcription unit be used, including a
30 pol III promoter, capping signal, and termination sequence. Pol III promoters, and other pol III transcription signals, are present in tRNA genes, 5S RNA genes, small nuclear RNA genes, and small cytoplasmic RNA

genes. Preferred pol III promoters for use in aptamer expression vectors are the human small nuclear U6 gene promoter and tRNA gene promoters. The use of U6 gene transcription signals to produce short RNA molecules *in vivo* is described by Noonberg *et al.*, *Nucleic Acids Res.* 22:2830-2836 (1995),
5 and the use of tRNA transcription signals is described by Thompson *et al.*, *Nucleic Acids Res.*, 23:2259-2268 (1995).

Many pol III promoters are internal, that is, they are within the transcription unit. Thus, these pol III transcripts include promoter sequences. To be useful for expression of aptamer molecules, these promoter sequences
10 should not interfere with the structure or function of the aptamer. The U6 gene promoter is not internal (Kunkel and Pederson, *Nucleic Acids Res.* 18:7371-7379 (1989); Kunkel *et al.*, *Proc. Natl. Acad. Sci. USA* 83:8575-8579 (1987); Reddy *et al.*, *J. Biol. Chem.* 262:75-81 (1987)). Suitable pol III promoter systems useful for expression of aptamer molecules
15 are described by Hall *et al.*, *Cell* 29:3-5 (1982), Nielsen *et al.*, *Nucleic Acids Res.* 21:3631-3636 (1993), Fowlkes and Shenk, *Cell* 22:405-413 (1980), Gupta and Reddy, *Nucleic Acids Res.* 19:2073-2075 (1990), Kickoefer *et al.*, *J. Biol. Chem.* 268:7868-7873 (1993), and Romero and Blackburn, *Cell* 67:343-353 (1991). The use of pol III promoters for expression of RNA
20 molecules is also described in WO 95/23225 by Ribozyme Pharmaceuticals, Inc.

In many contexts, it is preferred that expression of the aptamers is regulated, and in particular that the aptamers are expressed in a tissue- or cell-specific manner. For example, where a particular tissue or cell type is th
25 therapeutic focus, expression directed at that tissue or cell type is preferred. Many tissue- and cell-specific promoters are known and can be used for the expression of the disclosed aptamers. An example of this type of promoter would be the prostate specific antigen promoter, which is specifically activated in prostate cells and has been utilized for the targeting of anticancer
30 therapeutics of prostate tumors.

RNA fusion constructs can be produced which aid in the correct and specific targeting of expressed RNA molecules. These types of RNA fusion

constructs are discussed in detail in Good et al., "Expression of Small Therapeutic RNAs in Cell Nuclei", *Gene Therapy*, 4:45-54 (1997).

Expression vectors based on human tRNA(met) and U6 snRNA promoters are useful for targeting RNA expression and to render the resulting RNA

5 transcripts more resistant to degradation.

Other promoters useful for expressing the disclosed oligomers in mammalian host cells may be obtained from various sources, for example, the genomes of viruses such as: polyoma, Simian Virus 40 (SV40), adenovirus, retroviruses, hepatitis-B virus and most preferably

10 cytomegalovirus, or from heterologous mammalian promoters, e.g. beta actin promoter. The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment which also contains the SV40 viral origin of replication (Fiers et al., *Nature*, 273: 113 (1978)). The immediate early promoter of the human cytomegalovirus is conveniently obtained as a
15 HindIII E restriction fragment (Greenway, P.J. et al., *Gene* 18: 355-360 (1982)). Of course, promoters from the host cell or related species also are useful herein.

Enhancer generally refers to a sequence of DNA that functions at no fixed distance from the transcription start site and can be either 5' (Laimins,
20 L. et al., *Proc. Natl. Acad. Sci.* 78: 993 (1981)) or 3' (Lusky, M.L., et al., *Mol. Cell Bio.* 3: 1108 (1983)) to the transcription unit. Furthermore, enhancers can be within an intron (Banerji, J.L. et al., *Cell* 33: 729 (1983)) as well as within the coding sequence itself (Osborne, T.F., et al., *Mol. Cell Bio.* 4: 1293 (1984)). They are usually between 10 and 300 bp in length,
25 and they function in cis. Enhancers function to increase transcription from nearby promoters. Enhancers also often contain response elements that mediate the regulation of transcription. Promoters can also contain response elements that mediate the regulation of transcription. Enhancers often determine the regulation of expression of a gene. While many enhancer
30 sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein and insulin), typically one will use an enhancer from a eukaryotic cell virus. Preferred examples are the SV40 enhancer on the late

side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

5 The promotor and/or enhancer may be specifically activated either by light or specific chemical events which trigger their function. Systems can be regulated by reagents such as tetracycline and dexamethasone. There are also ways to enhance viral vector gene expression by exposure to irradiation, such as gamma irradiation, or alkylating chemotherapy drugs.

10 Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human or nucleated cells) may also contain sequences necessary for the termination of transcription which may affect mRNA expression. These regions are transcribed as polyadenylated segments in the untranslated portion of the mRNA encoding tissue factor protein. The 3' untranslated regions also include transcription termination sites. It is
15 preferred that the transcription unit also contain a polyadenylation region. One benefit of this region is that it increases the likelihood that the transcribed unit will be processed and transported like mRNA. The identification and use of polyadenylation signals in expression constructs is well established. It is preferred that homologous polyadenylation signals be
20 used in the transgene constructs. In a preferred embodiment of the transcription unit, the polyadenylation region is derived from the SV40 early polyadenylation signal and consists of about 400 bases (Figure 1). It is also preferred that the transcribed units contain other standard sequences alone or in combination with the above sequences improve expression from, or
25 stability of, the construct.

D. Capture Tags

Isolation of Ku:oligomer complexes can be facilitated through the use of capture tags coupled to the oligomer. As used herein, a capture tag is any compound that can be associated with a synthesized RNA molecule and which
30 can be used to separate compounds or complexes having the capture tag from those that do not. Preferably, a capture tag is a compound, such as a ligand or hapten, that binds to or interacts with another compound, such as a ligand-

binding molecule or an antibody. It is also preferred that such interaction between the capture tag and the capturing component be a specific interaction, such as between a hapten and an antibody or a ligand and a ligand-binding molecule.

5 Suitable capture tags include hapten or ligand molecules that can be coupled to the 5' end of the synthesized RNA molecule. Preferred capture tags, described in the context of nucleic acid probes, have been described by Syvanen *et al.*, *Nucleic Acids Res.*, **14**:5037 (1986)). Preferred capture tags include biotin, which can be incorporated into nucleic acids (Langer *et al.*,
10 *Proc. Natl. Acad. Sci. USA* **78**:6633 (1981)) and captured using streptavidin or biotin-specific antibodies. A preferred hapten for use as a capture tag is digoxigenin (Kerkhof, *Anal. Biochem.* **205**:359-364 (1992)). Many compounds for which a specific antibody is known or for which a specific antibody can be generated can be used as capture tags. Such capture tags can
15 be captured by antibodies which recognize the compound. Antibodies useful as capture tags can be obtained commercially or produced using well established methods. For example, Johnstone and Thorpe, *Immunochimistry In Practice* (Blackwell Scientific Publications, Oxford, England, 1987), on pages 30-85, describe general methods useful for producing both polyclonal and monoclonal
20 antibodies.

 Another preferred capture tag are anti-antibody antibodies. Such anti-antibody antibodies and their use are well known. For example, anti-antibody antibodies that are specific for antibodies of a certain class (for example, IgG, IgM), or antibodies of a certain species (for example, anti-rabbit antibodies) are
25 commonly used to detect or bind other groups of antibodies. Thus, one can have an antibody to the capture tag and then this antibody:capture tag:RNA complex can then be purified by binding to an antibody for the antibody portion of the complex.

 Another preferred capture tag is one which can form selectable
30 cleavable covalent bonds with other molecules of choice. For example, a preferred capture tag of this type is one which contains a sulfur atom. An RNA molecule which is associated with this capture tag can be purified by retention

on a thiolpropyl sepharose column. Extensive washing of the column removes unwanted molecules and reduction with β -mercaptoethanol, for example, allows the desired RNA molecules to be collected after purification under relatively gentle conditions (See Lorsch and Szostak, 1994 for a reduction to practice of this type of capture tag).

One preferred type of capture tag is a nucleotide cap analog, such as a 5'-5' guanosine dinucleotide cap. The capped aptamer would be incubated with a Ku-containing biochemical fraction, and a fusion protein containing the murine cap-binding protein would then be used to recover the aptamer-bound complex. This procedure has an advantage over antibody co-precipitation methods because the use of an aptamer that targets the DNA-binding site itself completely eliminates artifactual co-precipitation mediated by contaminating DNA, which is a major problem with co-precipitation of Ku protein. Bound proteins can be identified by SDS-PAGE, followed by other appropriate analytical methods.

Capture tags (and the associated oligomer) are isolated by interacting the capture tag with an appropriate molecule that binds the capture tag. These molecules, herein are referred to as capture tag receptors (CTRs). These CTRs can be associated with a solid support. When capture tag complexes are bound to CTRs attached to solid supports they can be effectively purified from the unwanted molecules because the solid support allows for successive washing to remove unbound molecules.

Supports that the CTRs can be coupled to can be any solid material to which the CTRs can be adhered or coupled. This includes materials such as acrylamide, cellulose, nitrocellulose, glass, polystyrene, polyethylene vinyl acetate, polypropylene, polymethacrylate, polyethylene, polyethylene oxide, glass, polysilicates, polycarbonates, teflon, fluorocarbons, nylon, silicon rubber, polyanhydrides, polyglycolic acid, polylactic acid, polyorthoesters, polypropylfumerate, collagen, glycosaminoglycans, and polyamino acids. Supports can have any useful form including thin films or membranes, beads, bottles, dishes, fibers, woven fibers, shaped polymers, particles and

microparticles. Preferred forms of supports are plates and beads. The most preferred form of beads are magnetic beads.

Methods for immobilization of oligonucleotides to substrates are well established. Oligonucleotides, including oligonucleotide capture docks, can be coupled to substrates using established coupling methods. For example, suitable attachment methods are described by Pease *et al.*, *Proc. Natl. Acad. Sci. USA* 91(11):5022-5026 (1994), and Khrapko *et al.*, *Mol Biol (Mosk) (USSR)* 25:718-730 (1991). A method for immobilization of 3'-amine oligonucleotides on casein-coated slides is described by Stimpson *et al.*, *Proc. Natl. Acad. Sci. USA* 92:6379-6383 (1995). A preferred method of attaching oligonucleotides to solid-state substrates is described by Guo *et al.*, *Nucleic Acids Res.* 22:5456-5465 (1994).

Methods for immobilizing antibodies to solid-state substrates are well established. Immobilization can be accomplished by attachment, for example, to aminated surfaces, carboxylated surfaces or hydroxylated surfaces using standard immobilization chemistries. Examples of attachment agents are cyanogen bromide, succinimide, aldehydes, tosyl chloride, avidin-biotin, photocrosslinkable agents, epoxides and maleimides. A preferred attachment agent is glutaraldehyde. These and other attachment agents, as well as methods for their use in attachment, are described in *Protein immobilization: fundamentals and applications*, Richard F. Taylor, ed. (M. Dekker, New York, 1991), Johnstone and Thorpe, *Immunochemistry In Practice* (Blackwell Scientific Publications, Oxford, England, 1987) pages 209-216 and 241-242, and *Immobilized Affinity Ligands*, Craig T. Hermanson *et al.*, eds. (Academic Press, New York, 1992). Antibodies can be attached to a support by chemically cross-linking a free amino group on the antibody to reactive side groups present within the solid-state support. For example, antibodies may be chemically cross-linked to a support that contains free amino or carboxyl groups using glutaraldehyde or carbodiimides as cross-linker agents. In this method, aqueous solutions containing free antibodies are incubated with the solid-state substrate in the presence of glutaraldehyde or carbodiimide. For crosslinking with glutaraldehyde the reactants can be incubated with 2%

glutaraldehyde by volume in a buffered solution such as 0.1 M sodium cacodylate at pH 7.4. Other standard immobilization chemistries are known by those of skill in the art.

In addition, non-antibody proteins such as streptavidin, can be linked using similar methods. Many protein and antibody columns are commercially available as well as specifically derivatized supports for conjugation to the CTRs.

E. Kits

The disclosed oligomers can also be used in a reagent kit. For example, kits for identification of the Ku protein or proteins that bind the Ku protein which contains, for example, an oligomer and suitable buffer substances. In this case the oligomer and the buffer substances can be present in the form of solutions, suspensions or solids such as powders or lyophilisates. The reagents can be present together, separated from one another or optionally also on a suitable carrier. The disclosed oligomers can also be used as a diagnostic agent or to identify the function of unknown genes.

II. Methods

The disclosed oligomers can be used for any purpose involving Ku binding to DNA, including methods to affect Ku activities and processes that depend on Ku activity and isolation of proteins that interact with Ku. For example, the disclosed oligomers can be used in pharmaceutical mixtures that contain one or several oligomers as the active substance, and, optionally, pharmaceutically acceptable auxiliary substances, additives and carriers. Such pharmaceutical oligomers are suitable for the production of an agent to specifically inhibit Ku DNA binding activity in cells or inhibit the activity of DNA-PK.

A. Method To Sensitize Cells To Agents

By introducing anti-Ku aptamers into cells, they can sensitize cells and tissues to the cytotoxic effects of therapeutic radiation and DNA-damaging drugs. This can be useful, for example, in cancer therapy where cancer cells can be made more sensitive to chemotherapeutic agents. Ku

protein and DNA-PK are important in the repair of radiation-induced DNA damage. If damage cannot be repaired, cells die. The cytotoxic effect of ionizing radiation forms the basis for radiation therapy, which is widely used in the treatment of human cancer. The efficacy of radiation therapy is currently limited by the radiation resistance of certain tumors (for example, glioblastomas) and by the side effects caused by irradiation of nearby normal tissues (for example, in treatment of breast and cervical cancer). Therefore, it is expected to be useful to have a means for sensitizing target cells and tissues to therapeutic radiation.

Inhibition of Ku functions can leave cells more sensitive to a variety of agents such as agents that damage DNA (discussed above) and agents that interfere with DNA topoisomerase function. For example, etoposides prevent cell division by interfering with DNA topoisomerase function. Jin et al., "Differential Etoposide Sensitivity of Cells Deficient in the Ku and DNA-PKcs Components of the DNA-dependent Protein Kinase" *Carcinogenesis* 19:965-97 (1998), describe the increased sensitivity of cells with Ku mutations to etoposides, which are inhibitors of DNA Topoisomerase II. Thus, a preferred therapeutic use of the disclosed oligomers is in combination with agents that render cells more sensitive to reduced Ku function. In this context, the aptamers can increase the effectiveness of the agent.

B. Identification Of Cellular RNA Sequences Predicted That Bind Ku Protein.

The disclosed aptamers are useful for identifying naturally occurring RNAs that bind to Ku protein. The existence of RNAs that bind tightly to Ku protein is consistent with suggestions that Ku protein function may be regulated to some extent by RNA in the normal cell. (Reeves, J. Exp. Med. 161, 18-39 (1985)). One site where such interactions might occur is in the nucleolus. The association of Ku protein with the nucleolus has been shown to be RNase sensitive (Reeves, J. Exp. Med. 161, 18-39 (1985)). A number of studies have shown that DNA-PK and Ku protein have an ability to regulate RNA polymerase I, which is localized in the nucleolus (Hoff et al.,

Proc. Natl. Acad. Sci. USA 91, 762-766 (1994); Kuhn et al., Genes and Development 9, 193-203 (1995); Labhart, Proc. Natl. Acad. Sci. USA 92, 2934-2938 (1995)). If, however, Ku protein associates with nucleolar RNAs, this would provide a potential targeting mechanism. Nucleoli contain a
5 multitude of discrete small nucleolar RNAs, as well as nascent ribosomal RNA (reviewed in Smith & Steitz, Cell 89, 669-672 (1997)). The consensus sequences present in the aptamers can be used to predict whether particular nucleolar RNAs or other cellular RNAs are likely to bind to Ku protein. Cellular RNA sequences that become available in public databases can be
10 searched to determine whether there is identity with the consensus sequences identified in the aptamers. Therefore, the disclosed aptamers are useful tools for predicting whether naturally occurring RNAs are likely to bind to Ku protein.

C. Identification Of Proteins That Interact With Ku Protein.

15 The disclosed aptamers can also be used to identify proteins that interact with Ku protein. This can be accomplished, for example, by attaching a chemical capture tag to the aptamer which allows the aptamer to be captured on a solid matrix. Ku aptamers attached to a chemical capture tag can be isolated on, for example, a solid support. The aptamer can then be
20 used to isolate Ku protein from a mixture. Any other molecules or proteins attached to the Ku protein can then be isolated also.

Another method to detect proteins that interact with Ku protein is to form complexes between radiolabeled aptamer and Ku protein, further incubate these complexes with the protein or mixture to be tested, and
25 analyze the complexes by native polyacrylamide gel electrophoresis. The aptamer-Ku protein complexes migrate as a discrete species at a defined position. The mobility of this species will be shifted and it will be found at a different position on the gel when additional proteins are bound.

The same assay can be adapted to detect proteins that promote
30 association of Ku protein and DNA-PKcs. Because the aptamers do not activate DNA-PK enzyme activity, it is expected that they will not, in themselves, promote assembly of Ku protein and DNA-PKcs into a

functional DNA-PK complex. That is, when purified DNA-PKcs are incubated with aptamer-Ku protein complexes, there should be no shift in mobility. This property can serve as an assay for proteins that promote association of Ku protein and DNA-PKcs. Complexes will be formed
5 between radiolabeled aptamer and Ku protein, and will be further incubated with DNA-PKcs and the protein or mixture to be tested. Complexes will be analyzed by native polyacrylamide gel electrophoresis. The mobility of the aptamer-Ku protein complex will be shifted when DNA-PKcs and additional proteins are bound. The presence of DNA-PKcs in the complex will be
10 ascertained by determining whether monoclonal antibodies against DNA-PKcs further shift the mobility.

Although a number of antibodies to Ku protein have been described, aptamers have different and complementary properties. A variety of commercially available monoclonal antibodies (N3H10, N9C1, 111, 162,
15 and S5C11) have been screened, as described in the examples, and all were found to produce supershifts in an electrophoretic mobility shift assay, indicating that they recognize epitopes outside the DNA binding site. By contrast, the aptamers competitively inhibited DNA binding, indicating that they are targeted directly to the DNA binding site of Ku protein. This
20 property makes the aptamers uniquely useful for probing the role of Ku protein in various physiological processes.

D. Manipulation Of Ku Activity In Cells.

The disclosed aptamers are also useful as tools for manipulation of Ku activity in cells. For this purpose, the aptamers should be expressed in or
25 delivered to cells. There are a number of techniques that can be used for targeted expression of small RNAs in the nuclear compartment (Sullenger et al., Mol. Cell. Biol. 10, 6512-6523 (1990); Thompson et al., Nucleic Acids Res. 23, 2259-2268 (1995); Good et al., Gene Therapy 4:45-54 (1997)). Introduction of aptamers to cells will be useful for investigations of the
30 physiological role of DNA-PK and other molecules and processes that Ku affects. Although DNA-PK is clearly involved in DNA repair, it has also been proposed that it is involved in a number of other processes, including

the stress response, viral infection, and transcriptional regulation (McConnell & Dynan, Curr. Opin. Cell Biol. 8, 325-330 (1996) and references therein).

A mechanism for manipulating DNA-PK activity *in vivo* is expected to provide a path for investigating the role of Ku protein in various

5 physiological processes.

E. Use Of Aptamers To Treat Autoimmune Disease.

The disclosed aptamers can also be used to treat autoimmune disease in patients producing anti-Ku antibodies. Antibodies to Ku protein occur in

15 15-20% of patients with systemic lupus erythematosus, scleroderma, myositis, and Sjogren's syndrome (Yaneva & Arentt, Clinical & Experimental Immunology 76, 366-372 (1989)). The immunogen in such cases may be a Ku protein-DNA complex, such as might be released from damaged or dying tissue. The disclosed aptamers can be used to disrupt such complexes, thus mitigating this chronic stimulus to the immune system.

20 For example, an aptamer may bind to Ku protein in a manner that does not allow an antibody to bind to Ku protein or may inhibit Ku protein binding to surface immunoglobulins on B cells. It has been suggested that self antigens are captured by antigen-specific receptors (surface immunoglobulins) on autoreactive B lymphocytes. (Craft and Fatenejad, *Arthritis and Rheumatism*, 40:1374-1382 (1997)). Thus, an aptamer that interferes with the binding of the Ku protein to the B cell, or which alters subsequent processing, is expected to block the chain of events leading to autoimmune disease.

F. Delivery Of Oligomers To Cells.

25 A variety of methods are available for delivering the disclosed oligomers to cells. For example, in general, the disclosed oligomers can be incorporated within or on microparticles or liposomes. Microparticles or liposomes containing the disclosed oligomers can be administered systemically, for example, by intravenous or intraperitoneal administration, 30 in an amount effective for delivery of the disclosed oligomers to targeted cells. Other possible routes include trans-dermal or oral, when used in conjunction with appropriate microparticles. Generally, the total amount of

the liposome-associated oligomer administered to an individual will be less than the amount of the unassociated oligomer that must be administered for the same desired or intended effect.

Oligomers including various polymers such as the polylactic acid and polyglycolic acid copolymers, polyethylene, and polyorthoesters and the disclosed oligomers can be delivered locally to the appropriate cells by using a catheter or syringe. Other means of delivering such oligomers locally to cells include using infusion pumps (for example, from Alza Corporation, Palo Alto, California) or incorporating the oligomers into polymeric implants (see, for example, Johnson and Lloyd-Jones, eds., *Drug Delivery Systems* (Chichester, England: Ellis Horwood Ltd., 1987), which can effect a sustained release of the therapeutic oligomers to the immediate area of the implant.

Delivery of the disclosed oligomers can be facilitated by targeting them to particular tissue or cell types. For example, the disclosed oligomers may be targeted to a particular cell type via antibodies, receptors, or receptor ligands. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Senter, et al., Bioconjugate Chem., 2:447-451, (1991); Bagshawe, K.D., Br. J. Cancer, 60:275-281, (1989); Bagshawe, et al., Br. J. Cancer, 58:700-703, (1988); Senter, et al., Bioconjugate Chem., 4:3-9, (1993); Battelli, et al., Cancer Immunol. Immunother., 35:421-425, (1992); Pietersz and McKenzie, Immunolog. Reviews, 129:57-80, (1992); and Roffler, et al., Biochem. Pharmacol., 42:2062-2065, (1991)). Vehicles such as "stealth" and other antibody conjugated liposomes (including lipid mediated drug targeting to colonic carcinoma), receptor mediated targeting of DNA through cell specific ligands, lymphocyte directed tumor targeting, and highly specific therapeutic retroviral targeting of murine glioma cells *in vivo*. The following references are examples of the use of this technology to target specific proteins to tumor tissue (Hughes et al., Cancer Research, 49:6214-6220, (1989); and Litzinger and Huang, Biochimica et Biophysica Acta, 1104:179-187, (1992)). In general, receptors are involved in pathways of endocytosis, either

constitutive or ligand induced. These receptors cluster in clathrin-coated pits, enter the cell via clathrin-coated vesicles, pass through an acidified endosome in which the receptors are sorted, and then either recycle to the cell surface, become stored intracellularly, or are degraded in lysosomes.

- 5 The internalization pathways serve a variety of functions, such as nutrient uptake, removal of activated proteins, clearance of macromolecules, opportunistic entry of viruses and toxins, dissociation and degradation of ligand, and receptor-level regulation. Many receptors follow more than one intracellular pathway, depending on the cell type, receptor concentration,
10 type of ligand, ligand valency, and ligand concentration. Molecular and cellular mechanisms of receptor-mediated endocytosis has been reviewed (Brown and Greene, DNA and Cell Biology 10:6, 399-409 (1991)).

- The oligomers can also be delivered to cells and expressed within the targeted cells using gene transfer technology based on viral and retroviral
15 vectors. Gene transfer can be obtained using direct transfer of genetic material, in a plasmid or viral vector, or via transfer of genetic material. Further, these methods can be used to target certain diseases and cell populations by using the targeting characteristics of the carrier. Transfer vectors can be any nucleotide construction used to deliver genes into cells
20 (e.g., a plasmid), or as part of a general strategy to deliver genes, e.g., as part of recombinant retrovirus or adenovirus (Ram et al. Cancer Res. 53:83-88, (1993)). Appropriate means for transfection, including viral vectors, chemical transfectants, or physico-mechanical methods such as electroporation and direct diffusion of DNA, are described by, for example,
25 Wolff, J. A., et al., Science, **247**, 1465-1468, (1990); and Wolff, J. A. Nature, **352**, 815-818, (1991).

III. Definitions

As used herein, the term "aptamer" is an oligomer or oligonucleotide that folds into a specific conformation and binds a target biomolecule.

- 30 As used herein, oligomer refers to oligomeric molecules composed of subunits where the subunits can be of the same class (such as nucleotides) or a mixture of classes (such as nucleotides and ethylene glycol). It is preferred

that the disclosed oligomers be oligomeric sequences, non-nucleotide linkers, or a combination of oligomeric sequences and non-nucleotide linkers. It is more preferred that the disclosed oligomers be oligomeric sequences.

Oligomeric sequences are oligomeric molecules where each of the subunits
5 includes a nucleobase (that is, the base portion of a nucleotide or nucleotide analogue) which can interact with other oligomeric sequences in a base-specific manner. The hybridization of nucleic acid strands is a preferred example of such base-specific interactions. Oligomeric sequences preferably are comprised of nucleotides, nucleotide analogues, or both, or are
10 oligonucleotide analogues.

Non-nucleotide linkers can be any molecule, which is not an oligomeric sequence, that can be covalently coupled to an oligomeric sequence. Preferred non-nucleotide linkers are oligomeric molecules formed of non-nucleotide subunits. Examples of such non-nucleotide linkers are
15 described by Letsinger and Wu, (*J. Am. Chem. Soc.* 117:7323-7328 (1995)), Benseler *et al.*, (*J. Am. Chem. Soc.* 115:8483-8484 (1993)) and Fu *et al.*, (*J. Am. Chem. Soc.* 116:4591-4598 (1994)). Preferred non-nucleotide linkers, or subunits for non-nucleotide linkers, include substituted or unsubstituted C₁-C₁₀ straight chain or branched alkyl, substituted or unsubstituted C₂-C₁₀
20 straight chain or branched alkenyl, substituted or unsubstituted C₂-C₁₀ straight chain or branched alkynyl, substituted or unsubstituted C₁-C₁₀ straight chain or branched alkoxy, substituted or unsubstituted C₂-C₁₀ straight chain or branched alkenyloxy, and substituted or unsubstituted C₂-C₁₀ straight chain or branched alkynyloxy. The substituents for these
25 preferred non-nucleotide linkers (or subunits) can be halogen, cyano, amino, carboxy, ester, ether, carboxamide, hydroxy, or mercapto.

As used herein, nucleoside refers to adenosine, guanosine, cytidine, uridine, 2'-deoxyadenosine, 2'-deoxyguanosine, 2'-deoxycytidine, or thymidine. A nucleoside analogue is a chemically modified form of
30 nucleoside containing a chemical modification at any position on the base or sugar portion of the nucleoside. As used herein, the term nucleoside analogue encompasses, for example, both nucleoside analogues based on

naturally occurring modified nucleosides, such as inosine and pseudouridine, and nucleoside analogues having other modifications, such as modifications to the 2' position of the sugar. As used herein, nucleotide refers to a phosphate derivative of nucleosides as described above, and a nucleotide
5 analogue is a phosphate derivative of nucleoside analogues as described above. The subunits of oligonucleotide analogues, such as peptide nucleic acids, are also considered to be nucleotide analogues.

As used herein, a ribonucleotide is a nucleotide having a 2' hydroxyl function. Analogously, a 2'-deoxyribonucleotide is a nucleotide having only
10 2' hydrogens. Thus, ribonucleotides and deoxyribonucleotides as used herein refer to naturally occurring nucleotides having nucleoside components adenosine, guanosine, cytidine, and uridine, or 2'-deoxyadenosine, 2'-deoxyguanosine, 2'-deoxycytidine, and thymidine, respectively, without any chemical modification. Ribonucleosides, deoxyribonucleosides,
15 ribonucleoside analogues and deoxyribonucleoside analogues are similarly defined except that they lack the phosphate group, or an analogue of the phosphate group, found in nucleotides and nucleotide analogues.

As used herein, oligonucleotide analogues are polymers of nucleic acid-like material with nucleic acid-like properties, such as sequence
20 dependent hybridization, that contain at one or more positions, a modification away from a standard RNA or DNA nucleotide. A preferred example of an oligonucleotide analogue is peptide nucleic acid.

As used herein, base pair refers to a pair of nucleotides or nucleotide analogues which interact through one or more hydrogen bonds. The term
25 base pair is not limited to interactions generally characterized as Watson-Crick base pairs, but includes non-canonical or sheared base pair interactions (Topal and Fresco, *Nature* 263:285 (1976); Lomant and Fresco, *Prog. Nucl. Acid Res. Mol. Biol.* 15:185 (1975)).

As used herein the term "gene" when used in the context of genetic
30 vectors refers to the genetic material of interest that is to be transferred to the host organism. For example, the "gene" to be transferred could be a

sequence which codes for an aptamer and when transcribed would produce a functional aptamer.

The present invention will be further understood by reference to the following examples.

5

Examples

Example 1: Preparing the RNA Pool and Screening for Ku protein binding.

Because there was little previous information about Ku protein-RNA interactions, the RNA binding properties of Ku protein were systematically investigated using SELEX (systematic evolution of ligands by exponential enrichment) technology. With this technology, it is possible to identify oligonucleotides from a large, random pool that bind to a ligand of interest. The SELEX method has been used to identify oligonucleotides that bind to nucleic acid binding proteins, to non-nucleic acid binding proteins, and to small molecules (reviewed in Gold et al., Annu. Rev. Biochem. 64, 763-797 (1995)).

The RNA pool was prepared by synthesis of single stranded RNA from a template comprising a region of conserved sequences and a region of randomized and/or biased sequences.

20 Synthesis of the RNA Pool

SELEX protocols and the template used to synthesize the nonselected RNA pool were obtained from Dr. Hang Chen and Dr. Larry Gold (University of Colorado) and modified as described below. The major change to the SELEX protocol was to increase the KCl concentration during sequential rounds of SELEX to maintain stringency. Protocols can also be obtained from Tuerk & Gold, Science 249, 505-510 (1990), and U.S. Patent Numbers 5,270,163 and 5,567,588. HeLa cell nuclear extracts and oligonucleotides were obtained from Dr. Mark Anderson and Ms. Kirsten Strickler. Chemicals were obtained from Sigma, Fisher Scientific, antibodies from Neomarkers, nitrocellulose filters from Millipore, cloning vectors from Promega and Invitrogen, translation kits from Ambion, and RNasin from Promega.

SELEX procedures to select Ku binding RNAs were carried out largely as described in Chen & Gold, *Biochemistry* 33, 8746-8756 (1994). The template for synthesis of the starting RNA pool was based on a DNA oligonucleotide, 5'-CCCGGATCCTAGTTCACGATGCTGCAA-(N)₄₀-TTACGGTCTGAGAAAATATCCTCCC-3', where N indicates an equimolar mixture of A, G, C and T. This sequence was used because it is known to function well in SELEX. The 5' and 3' constant regions do not bind to Ku protein. DNA templates were generated by PCR amplification of this oligonucleotide using primer 1, 5'-CCCAAGCTTAATAC GACTCACTATAGGGAGGATATTTTC TCAGACCGTAA-3', and primer 2, 5'-CCCGGATCCTAGTTCACGATGC TGCAA-3'. Amplification was carried out with 0.5 μ M of the oligonucleotide containing the random segment, 2 μ M primer 1 and primer 2, 1 mM dATP, dCTP, dGTP, and dTTP, 50 U/ml Taq DNA polymerase, 50 mM KCl, 10 mM Tris-HCl, pH 8.4, 7.5 mM MgCl₂, and 50 μ g/mL BSA in a volume of 0.5 ml. After four cycles of amplification (93° C for 30 s, 57° C for 20 s, 73° C for 90 s), the reaction was extracted with an equal volume of phenol-chloroform-isoamyl alcohol (25:24:1 v/v/v) (PCIA), and nucleic acids were precipitated from the supernatant by addition of 0.1 vol of 3 M NaOAc pH 5.2 and 2.5 vol ethanol.

This template preparation was incubated for 2-3 h at 37° C in a reaction mixture containing 475 U/ml T7 RNA polymerase, 40 mM Tris-HCl, pH 8.0, 12 mM MgCl₂, 5 mM DTT, 1 mM spermidine, 0.002 % Triton X-100, 4 % PEG 8000, 2 mM ATP, UTP, and GTP, and 0.2 mM [α -³²P] CTP (1 Ci/mmol) in a final volume of 200 μ l. DNase I was added (25 U/ml) and incubation was continued for 30 minute. The resulting RNAs were fractionated by 10% urea-PAGE and gel slices containing full-length 92 nucleotide RNA were excised, crushed, mixed with 0.5 ml TE, 0.5 ml phenol, and 10 μ l 10% sodium dodecyl sulfate, and tumbled overnight at 4° C to extract RNA. The supernatant was collected and RNA was precipitated with NaOAc and EtOH as described above. RNA was dissolved in TE, denatured at 100° C for 2 minute, and renatured by adjusting to 5 mM

MgCl₂, cooling rapidly to 0° C, and incubating for 30 minute. The amount of RNA recovered was estimated by liquid scintillation counting.

Protein purification

Native DNA-PKcs and Ku protein were purified from HeLa cell
5 nuclear extracts as previously described in Dvir et al., J. Biol. Chem
268(14):10440-10447 (1993), except that the Phenyl-Superose and Mono S
steps were omitted. Monoclonal antibodies to Ku protein were obtained from
Neomarkers (Fremont, CA).

Selection of Ku-binding RNAs

10 To perform *in vitro* selections, the RNA pool was mixed with
purified Ku protein under conditions for complex binding. RNA and Ku
protein were incubated for 30 minute at room temperature in a buffer
containing 25 mM Tris-HCl, pH 7.9, 0.5 mM EDTA, 10 % glycerol, 5 mM
MgCl₂, 0.5 mM DTT, and 0.01 % Tween 20 in a final volume of 50-100 µl.
15 Concentrations of RNA, Ku protein, and KCl were variable, as specified in
Table 2. Ku protein-RNA complexes were isolated by electrophoresis on a 5
% non-denaturing polyacrylamide gel containing 25 mM Tris-HCl, pH 8.3,
190 mM glycine, and 1 mM EDTA at 300 V for one hour. In one
experiment, complexes were isolated by passage over a nitrocellulose filter
20 (Millipore 25-µm HAWP), which was washed twice with 5 ml of 25 mM
Tris-HCl, pH 7.9. RNA was eluted from the polyacrylamide gel or the filter
by tumbling in a mixture of TE, phenol, and SDS as described above.

The RNAs having a high affinity for the Ku protein relative to the
mixture were partitioned by nitrocellulose filter binding and electrophoretic
25 mobility shift assay. Another way to separate complexes is by
immunoprecipitation.

Once the enriched RNA molecules were separated from the isolated
complexes, they were amplified by reverse transcription followed by
polymerase chain reaction. The complexity of a selected pool can be
30 characterized by direct sequencing or restriction enzyme analysis.

Additional Rounds of SELEX Selection and Amplification

When additional rounds of selection were performed, purified Ku protein (from HeLa cell nuclear extracts) was mixed with the RNA pool and allowed to bind. Ku-RNA complexes were isolated using either a
5 nitrocellulose filter binding or an electrophoretic mobility shift assay (EMSA). RNA that was bound to Ku protein was isolated, amplified by RT-PCR, and again enriched based on the ability to bind to Ku protein. The selected RNA was reverse-transcribed by incubating in a reaction mixture containing 50 mM Tris-HCl, pH 8.3, 60 mM NaCl, 6 mM Mg(OAc)₂, 10
10 mM DTT, 0.4 mM dATP, dCTP, dGTP, and dTTP, and 300 U/ml of AMV reverse transcriptase, in a final volume of 50 μ l. The resulting cDNA was amplified by 15 cycles of PCR using primers 1 and 2 at 93°C for 30 seconds, 58°C for 20 seconds, 72°C for 90 seconds. This process was repeated 6-7 times. Selection conditions used for each round of selection are given in
15 Table 2. To maintain the stringency of selection, the RNA:protein ratio, the KCl concentration, or both were progressively increased as shown in Table 2. Two independent selections were conducted using different batches of starting RNA.

After six or seven rounds of selection, cDNA was synthesized from
20 the final RNA pool and amplified by 5 cycles of PCR using primer 1 and 2. The product was digested with Bam HI and Hind III and subcloned into the Bam HI and Hind III sites of pGEM3zf(+) (Promega) or pZero2.1 (Invitrogen). Inserts were sequenced with an m13 reverse primer using an ABI automated sequencer.

Table 2: Summary of the Selection Parameters

Selection I				
SELEX Round	Input RNA (nM)	Ku protein ^a (nM)	KCl ^b (mM)	Selection ^c Method
1	1305	64.1	50	NCFA
2	612.5	24.3	50	EMSA
3	860	42.6	120	EMSA
4	1130	42	120	EMSA
5	2320	64	150	EMSA
6	142	3.2	160	EMSA

Selection II				
SELEX Round	Input RNA (nM)	Ku protein (nM)	KCl (mM)	Selection Method
1	1330	53	120	EMSA
2	1660	42	120	EMSA
3	456	14.9	150	EMSA
4	144	3.84	160	EMSA
5	1065	21.3	160	EMSA
6	2710	18.6	160	EMSA
7	436	2.4	160	EMSA

^a The ratio of Ku protein to input RNA was gradually decreased in order to increase the stringency. ^b Salt concentration was increased to maintain high stringency. ^c Nitrocellulose filter binding assay (NCFA) or electrophoretic mobility shift assay (EMSA) were employed for selection methods.

5 RNAs from the last round of selection were cloned and sequenced using standard techniques. Table 1 presents the sequence of nineteen aptamer RNAs that were isolated that bind avidly to Ku protein. Many of the RNAs contain common sequence motifs. A number of the RNAs bind competitively with DNA to Ku protein and thereby inhibit DNA and PK
10 enzymatic activity. The ability of Ku protein to bind tightly to specific RNAs is consistent with a role for RNA in the regulation of Ku protein activity or intranuclear localization. Additionally, the identification of diverse RNAs that bind avidly to Ku protein raises the possibility that natural RNAs modulate the activity of DNA-PK *in vivo*.

15 Binding of Ku Protein

 Binding of RNA and DNA to Ku protein was measured using EMSA as shown by the representative Ku protein-RNA binding curves after successive rounds of SELEX in Figure 1. Binding curves were obtained with EMSA followed by a nonlinear least squares fit of the data. The RNA
20 pools in Figure 1 were transcribed from templates prior to the first round (nonselected), and after the fourth (4th), and the sixth (6th) rounds of SELEX using T7 RNA polymerase as described in example 2. A twenty-one base pair double-stranded DNA was end-labeled with T4 polynucleotide kinase. The DNA was arbitrarily chosen as a transcription factor binding site from
25 the human T-cell leukemia virus proviral promoter, 5'-CTCAGGCGTTGACGACAACCC-3'.

 Ku protein bound to nonselected RNA with an apparent average K_d of 24 nM. This binding was somewhat stronger than expected, given that previous work had found that Ku protein had little ability to interact with
30 tRNA or synthetic RNA polymers (Mimori & Hardin, J. Biol. Chem. 261, 10375-10379 (1986)). Pooled RNA tested after the 4th and 6th round of

SELEX showed an increased ability to bind to Ku protein. RNA obtained after the 6th round of selection bound with an apparent K_d of about 0.3 nM, which was comparable to the binding seen with a double-stranded DNA oligonucleotide under similar conditions. Although the average affinity of the selected RNA was only 75-fold greater than for the nonselected pool, subsequent experiments revealed clear functional differences, as only the selected RNAs were able to inhibit biochemical activities of Ku protein.

Example 2: Comparison of binding of RNAs to determine specificity.

Synthesis of aptamer RNAs

Individual aptamer RNAs were synthesized by T7 RNA polymerase using linearized plasmid template. Both radiolabeled RNA and nonradiolabeled RNA were prepared using a MEGA shortscript T7 kit (Ambion) with 8 μ g linearized plasmid template. The RNA was gel-purified, heated, and refolded as described in example 1, and the final concentration was determined spectrophotometrically. Secondary structures were predicted using the method of Zuker (Jaeger et al., Proc. Natl. Acad. Sci. USA 86, 7706-7710 (1989); Jaeger et al., Meth. Enzymol. 183, 281-306 (1990)) as implemented on the mfold server www.ibc.wustl.edu/%7Ezucker/rna/form1.cgi.

Synthesis of RNA containing the TAR sequence

Pools of RNA at various stages of selection were also characterized directly for their ability to bind to Ku protein in an electrophoretic mobility shift assay. For comparison of the aptamer RNAs binding affinity with other known RNAs, these assays also included a HIV TAR RNA, which was synthesized by T7 RNA polymerase as described in the preceding section. This RNA consisted of the following HIV-derived sequence:
GGGUCUCUCUGGUUAGACCAGAUUCUGAGCCUGGGAGCUCUCUG
GCUAACUAGGGAACCC.

TAR-containing RNAs have previously been reported to bind selectively to Ku protein (Kaczmarzski & Khan, Biochem. Biophys. Res. Commun. 196, 935-942 (1993)).

Binding of the TAR RNA was measurable, but weak, under the conditions of the experiments. These results confirm that TAR-containing RNA has some ability to bind to Ku protein, but indicate that the binding is not particularly strong, relative to other short RNAs. An *in vitro* yeast tRNA
5 transcript also bound very weakly to Ku protein.

Example 3: Cloning, sequence analysis of aptamers, and K_d of aptamers.

After six and seven rounds of SELEX (Selection I and II in Table 2, respectively), the pooled RNA was reverse-transcribed, PCR-amplified, and
10 cloned into plasmid vectors. 82 clones were isolated, including 63 from Selection I and 19 from Selection II. Allowing for duplicates, the 82 clones represented 19 independent sequences, which are given in Table 1.

An equilibrium dissociation constant was determined for the binding of each of the 19 different RNAs to Ku protein. The results, given in Table
15 1, showed that most of the RNAs have a K_d of less than 2 nM.

The sequence alignment in Table 1 revealed that the RNAs can be grouped into three classes, based on the presence of conserved sequence motifs in the variable region of each aptamer. The most striking of these motifs was a sequence, GCUUCCCCANNNAC (herein designated SEQ ID
20 NO:20) (wherein N may be A, C, G or T), which was perfectly conserved in three independent RNA sequences and partially conserved in four others. A second motif, AMAUGA (herein designated SEQ ID NO:21) (wherein M may be either A or C) was perfectly conserved in three RNAs and partially conserved in seven others. A third motif, which partially overlaps the
25 second, had the sequence AACUUCGA (bases 31-38 of SEQ ID NO:15). This motif was present in one RNA and partially conserved in four others. Several RNAs fell into hybrid classes containing two of the three motifs. Subsequent analysis showed that the hybrid RNAs containing two motifs were among the most efficacious blockers of Ku protein activity in
30 functional assays.

Example 4: Inhibition of DNA-PK activity by different RNA aptamers.

Binding of the RNA aptamers to Ku protein to determine if they are capable of regulating Ku protein activity was then measured. One of the major biochemical functions of Ku protein is to regulate the activity of DNA-PK. Ku protein targets the catalytic subunit of DNA-PK (DNA-PKcs) to DNA, increasing phosphorylation activity by 5-50 fold (Dvir et al., Proc. Natl. Acad. Sci. USA 89, 11920-11924 (1992); Gottlieb & Jackson, Cell 72, 131-142 (1993)). Under some assay conditions, DNA-PK that has been depleted of Ku protein can no longer be stimulated by double-stranded DNA (Dvir et al., J. Biol. Chem. 268, 10440-10447 (1993)).

Kinase assay

Peptide phosphorylation was carried out using a previously described method (Lees-Miller et al., Mol. Cell. Biol. 12, 5041-5049 (1992)) with modifications. Reactions contained 1 nM Ku protein, 0.5 nM DNA-PKcs, 100 μ M p53-derived peptide (EPPLSQEAFADLLWKK, phosphorylation site underlined), 0.25 nM DNA fragment (308 base pair Bgl I-Blp I fragment from pHSE1 (Peterson et al., J. Biol. Chem. 270, 1449-1454 (1995b)), 4000 U/ml RNasin (Promega), 0.2 mM [g - 32 P] ATP (1.25 Ci/mmol), 25 mM Tris-HCl, pH 7.9, 25 mM MgCl₂, 1.5 mM DTT, 50 mM KCl, and 10% glycerol in a total volume of 20 μ l. Reactions were incubated at 30° C for 45 minute, stopped by addition of 10 μ l of 1 mg/ml BSA and 10 μ l of 40 % TCA, and further incubated for 30 minute on ice. Precipitated protein was removed by centrifugation and 10 μ l of each supernatant, which contained the phosphorylated p53 peptide, was spotted on a phosphocellulose filter (Whatman P81). The filters were washed three times with 15% HOAc for 15 minutes, and incorporation of radiolabel was measured by liquid scintillation counting.

Ku-binding RNAs were able to block the ability of Ku protein to activate DNA-PK. Each of the nineteen selected aptamers, as well as the nonselected RNA pool, was tested in a DNA-PK assay. Each RNA was

prepared by *in vitro* transcription and added to a DNA-PK phosphorylation assay.

DNA-PK assay reactions contained 0.5 nM Ku protein, 1 nM DNA-PKcs, 0.25 nM 308-base pair double-stranded DNA fragment, a synthetic peptide (EPPLSQEAFADLLWKK) containing a DNA-PK phosphorylation site from p53 (underlined), and the indicated amounts (8, 16, 32 nM) of RNA aptamer. Native DNA-PKcs were purified from HeLa cell nuclear extracts as described in example 2. In Figure 2A, DNA-PK activity is expressed as a percentage of activity in the absence of RNA. Values shown are averages of duplicate reactions with standard deviations indicated. Background phosphorylation in the absence of DNA has been subtracted.

The effect on DNA-PK activity was measured, relative to control reactions with no added RNA. Nonselected RNA had no effect on DNA-PK activity (Figure 2A). Of the selected RNAs, seven inhibited DNA-PK activity by 85% or more when present at 16-32 nM (Figure 2A). Ten RNAs inhibited DNA-PK to a lesser extent and two had no effect on DNA-PK in the concentration range tested. These results indicate that at least some of the selected RNAs interact with regions of Ku protein that are critical for its biochemical activity. Since DNA-PKcs are known to have a low level of activity in the absence of Ku protein (Dvir et al., J. Biol. Chem. 268, 10440-10447 (1993)), the inhibition seen with the more effective RNAs probably reflects near-total ablation of Ku protein regulatory function.

RNAs cannot activate DNA-PK directly

It was determined whether any of the RNAs were capable of activating, as opposed to inhibiting, DNA-PK phosphorylation activity. None of the RNAs significantly increased DNA-PK activity in a standard assay performed in the presence of DNA. However, under these conditions, the DNA-PK was already highly active and a weak ability of RNA to activate DNA-PK might have gone undetected. Indeed, displacement of DNA by a weakly activating RNA may have resulted in partial inhibition of DNA-PK activity.

To address the question whether any of the RNAs were capable of activating DNA-PK in the absence of DNA, DNA-PK assays in the absence of DNA were performed. Kinase assays were performed as described above in the presence or absence of 0.25 nM double-stranded DNA, as indicated.

- 5 Reactions contained various amounts of aptamer RNA (8, 16, 32 nM).

When neither RNA nor DNA was present, DNA-PK had a low level of basal activity, as expected. The addition of double-stranded DNA increased this activity about ten-fold. Four RNAs were tested, including one that had given no effect on DNA-PK activity in the earlier assays (#52), two
10 that had given partial inhibition (SC1, SC9), and one that had given potent inhibition (SC4). None of these four RNAs showed any ability to activate DNA-PK above background levels in the absence of DNA.

Competitive binding analysis between double stranded DNA and RNA In principle, an RNA that binds to Ku protein could inhibit DNA-PK
15 either by blocking the binding of Ku protein to DNA or by blocking the interaction between Ku protein and DNA-PKcs. To distinguish these possibilities, the RNAs were tested for their effect on Ku protein-DNA binding.

Ku protein was incubated with a radiolabeled twenty-one base pair
20 double stranded DNA (1 nM) oligonucleotide in the presence or absence of various amounts (8, 16, 32 nM) of nonradiolabeled RNA. The incubation conditions were 25 mM Tris-HCl pH 7.9, 0.5 mM EDTA, 10% glycerol, 5 mM MgCl₂, 0.5 mM DTT, 0.01% Tween, 120 mM KCl, 30 minute at ambient temperature. Ku protein-DNA complexes were trapped in a
25 nitrocellulose filter binding assay as described in example 1 to measure Ku-DNA complexes. The results are shown in Figure 2B, where binding is expressed as a percentage of Ku-DNA complexes detected in the absence of competitor. Values shown are averages of duplicate reactions with standard deviations indicated. Eleven cloned RNAs were tested, all of which had
30 previously been shown to inhibit DNA-PK in a phosphorylation assay. All of these RNAs competed to some extent with the double-stranded DNA oligonucleotide for binding to Ku protein. The seven RNAs that functioned

as effective inhibitors of DNA-PK enzyme activity also proved to be effective inhibitors of Ku protein-DNA binding activity (Figure 2B). By contrast, RNA from the nonselected pool had no effect on Ku protein-DNA interaction (Figure 2B). There was a rough correlation between the efficiency with which individual RNAs inhibited DNA binding and DNA-PK phosphorylation. Although not wishing to be bound by theory, the results suggest that all of the inhibitory RNAs that were tested work primarily by interfering with protein-DNA interactions, rather than protein-protein interactions.

Both of the subunits of Ku protein have a predicted net negative charge at neutral pH, and it may be that unfavorable electrostatic interactions restrict interaction with aptamer RNAs over much of the protein's surface. Given this constraint, it is perhaps not surprising that many of the aptamers are targeted toward the DNA binding site within the Ku protein. Although this site has not been fully mapped, it is likely that it contains basic residues that are capable of electrostatic interactions with the phosphate backbone of RNA and DNA.

To further explore the relationship between RNA and DNA binding sites in Ku protein, Electrophoretic Mobility Shift Assays (EMSAs) were performed comparing radiolabeled aptamer RNA (1 nM) and radiolabeled twenty-one base pair double stranded DNA (1 nM) probes. Cross-competition between RNA aptamers and DNA for Ku protein binding were found. It was discovered that Ku protein forms stable complexes with both types of probes, and that the complexes with RNA and DNA have similar electrophoretic mobilities. Both of the RNAs that were tested, #7-3 and #SC9, competed with DNA for binding to Ku protein. RNA #7-3 is a more effective competitor than RNA #SC9, consistent with the results of nitrocellulose filter binding assays. In the reciprocal experiment, nonradiolabeled DNA fragments competed with labeled RNA for binding to Ku protein. As controls, DNA and each RNA were shown to effectively self-compete for binding to Ku protein, confirming the fidelity of the analysis.

Notably, neither the DNA nor the RNA competitors induced the formation of supershifted complexes. The absence of supershifted complexes provides additional evidence that RNA and DNA cannot bind simultaneously to Ku protein, but rather, bind competitively to the same site.

5 Specific inhibition of DNA-PK by aptamers in crude cell extracts

Part of the rationale for identifying RNAs that interact with Ku protein was to develop potential tools, aptamers, for *in vivo* modulation of DNA-PK activity. To be useful, such aptamers must not only recognize Ku protein, but they must be able to do so in a complex milieu containing other
10 macromolecules.

To determine whether the RNAs were specific for Ku protein, the ability of the aptamers to inhibit DNA-PK in crude HeLa cell nuclear extracts were measured. As shown in Figure 3, phosphorylation of the p53-derived peptide substrate by crude extracts is stimulated at least five-fold by
15 addition of double stranded DNA. This suggests that DNA-PK is the major kinase that phosphorylates this substrate under the conditions used. The reactions contained 2 μ l HeLa cell nuclear extract (0.4 μ g protein) and other components as in Figure 2. Reactions were performed in the presence of various amounts of aptamer RNA (250, 500, 1000 nM) as indicated. The
20 assays were as described above except a higher amount of DNA was used, 5.3 nM. 100 to 142.1 μ M p53 was used. DNA-PK activity is expressed as a percentage of activity in the absence of RNA. Values shown are averages of duplicate reactions with standard deviations as indicated. Four different aptamer RNAs, all of which had previously been shown to inhibit purified
25 DNA-PK, were tested in the crude extract. All four reduced enzyme activity to near-basal levels. RNA from the nonselected pool had no effect at concentrations up to 1000 nM (Figure 3, lanes 5-7). These results demonstrate that the aptamers are sufficiently selective that they can recognize Ku protein and inhibit DNA-PK activity even in a crude
30 preparation.

Dissociation of Ku-DNA complexes by aptamer RNAs

Previous work has suggested that, once the Ku protein binds to double-stranded DNA, the resulting complexes are not readily disrupted by a second DNA added as competitor, except when the first and second DNAs have cohesive ends (Bliss & Lane, J. Biol. Chem. 272, 5765-5773 (1997)). To find out whether aptamers were capable of capturing Ku protein from preformed complexes, an experiment was performed in which purified Ku protein (0.32 nM) was incubated with a radiolabeled double-stranded DNA (5' end labelled with T-4 polynucleotide kinase and [γ -32P] ATP) fragment (0.38 nM) under conditions for forming complexes for 30 minutes at room temperature. The complexes were challenged with either aptamer RNA or RNA from the nonselected pool. Incubation was then continued and the amount of Ku-DNA complexes remaining were measured by electrophoretic mobility assay using a 5% nondenaturing polyacrylamide gel containing 25 mM Tris-HCl pH 8.3, 190 mM glycine, 1 mM EDTA, run for one hour at 300 V, ambient temperature.

Ku protein-DNA complexes steadily dissociated in the presence of aptamer #7-3. In contrast, the amount of Ku protein-DNA complexes remained almost unchanged when challenged with nonselected RNA. This result demonstrates that the aptamer RNA can compete with double-stranded DNA for the binding of Ku protein even if the aptamer RNA is added after Ku-DNA complexes have formed.

Claims

We claim:

1. An oligomer comprising:
a sequence that specifically binds to Ku protein with an equilibrium dissociation constant less than 10 nM.
2. The oligomers according to claim 1 wherein the oligomers compete with DNA to bind Ku protein.
3. The oligomers according to claim 1 wherein the oligomer comprises at least one of the three nucleotide sequences chosen from the group comprising: GCUUCCCCANNAC, A(A/C)AUGA, and AACUUCGA.
4. An oligomer according to claim 1 wherein the oligomer is made of RNA nucleotides.
5. An oligomer according to claim 1 wherein the oligomer comprises the components A, A', B, C, C', D, and E, wherein,
the components are arranged 5'-A-B-C-D-C'-E-A'-3' wherein,
component A hybridizes to component A',
component C hybridizes to component C',
components B and E form bulges, and
component D forms a structure which is either a bulge or a loop.
6. An oligomer according to claim 5 wherein the secondary structure formed is represented by Figure 6A.
7. An oligomer comprising the sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, and SEQ ID NO:19.
8. A method of altering an activity of Ku protein comprising bringing into contact an oligomer of claim 1 and Ku protein.
9. The method of claim 8 wherein the Ku protein activity is the effect of Ku on DNA-PK.

10. The method of claim 8 wherein the bringing into contact is accomplished by delivering the oligomer to a cell or by expressing the oligomer in a cell.

11. The method of claim 10 wherein the cell becomes more sensitive to an agent when the oligomer is delivered to or expressed in the cell.

12. The method of claim 11 wherein the agent is a mutagen.

13. The method of claim 12 wherein the mutagen is radiation.

14. The method of claim 11 wherein the agent is a topoisomerase inhibitor.



15. A method of identifying physiological effects of Ku protein comprising exposing a cell to an oligomer of claim 1, and identifying a physiological difference between the cell exposed to the oligomer and a cell not exposed to the oligomer.

16. A method of isolating proteins that interact with Ku protein comprising

incubating an oligomer of claim 1 with a sample containing Ku protein such that oligomer-Ku complexes can form,
recovering the oligomer-Ku complexes, and
recovering any protein associated with the oligomer-Ku complexes.

17. The method of claim 16 wherein the oligomer further comprises a capture tag, and wherein the oligomer-Ku complex is recovered using the capture tag.

1/5

**FIG. 1****FIG. 2A**

SUBSTITUTE SHEET (RULE 26)



**FIG. 2B****FIG. 3**

FIG. 5






FIG. 6A**FIG. 6B**

INTERNATIONAL SEARCH REPORT

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/11 C12Q1/68 C07H21/02		International Application No PCT/US 98/27728
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N C12Q		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KACZMARSKI W ET AL: "LUPUS AUTOANTIGEN KU PROTEIN BINDS HIV-1 TAR RNA IN VITRO" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 196, no. 2, pages 935-942, XP002061470 ISSN: 0006-291X cited in the application the whole document <div style="text-align: center;">--- -/-</div>	1,2,4,5
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
* Special categories of cited documents: <div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">7 June 1999</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">17/06/1999</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Andres, S</div>

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/27728

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>GIFFIN, W. ET AL.: "Sequence-specific DNA binding and transcription factor phosphorylation by Ku autoantigen/DNA-dependent protein kinase" JOURNAL OF BIOLOGICAL CHEMISTRY., vol. 272, 28 February 1997 (1997-02-28), pages 5647-5658, XP002105043 cited in the application page 5650, right-hand column, line 15 -, sentence 48 page 5652, left-hand column, line 4 -, sentence 19 page 5655, right-hand column, line 17 -, sentence 20</p>	1,2,8,9
X	<p>LUDWIG D L ET AL: "Ku80 gene expression is Sp1-dependent and sensitive to CpG methylation within a novel cis element" GENE: AN INTERNATIONAL JOURNAL ON GENES AND GENOMES, vol. 199, no. 1-2, 15 October 1997 (1997-10-15), page 181-194 XP004126380 page 188, paragraph 3.5 - page 189, paragraph 3.6</p>	1,2,8
A	<p>HE, D. ET AL.: "Restoration of X-ray and etoposide resistance, Ku-end binding activity and V(D)J recombination to the chinese hamster xci-3 mutant by hamster Ku86 cDNA" MUTATION RESEARCH, vol. 363, 1996, pages 43-56, XP002105044 AMSTERDAM, NL ISSN: 0027-5107 the whole article, especially last sentence of paragraph 4.3 on page 53</p>	10-14
A	<p>WO 91 19813 A (UNIV COLORADO FOUNDATION) 26 December 1991 (1991-12-26) cited in the application page 8, line 33 - page 12, line 4 page 54, line 24 - page 56, line 28 claims figure 22</p>	
A	<p>BLISS, T. & LANE, D.: "Ku selectively transfers between DNA molecules with homologous ends" JOURNAL OF BIOLOGICAL CHEMISTRY., vol. 272, 28 February 1997 (1997-02-28), pages 5765-5773, XP002105045 cited in the application</p>	

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/27728

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	YOO S ET AL: "Characterization of the RNA binding properties of Ku protein." BIOCHEMISTRY, (1998 FEB 3) 37 (5) 1336-43., XP002105046 the whole document ----	1-9
T	DYNAN W S ET AL: "Interaction of Ku protein and DNA - dependent protein kinase catalytic subunit with nucleic acids." NUCLEIC ACIDS RESEARCH, (1998 APR 1) 26 (7) 1551-9., XP002105047 -----	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/27728

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 8 to 14 (as far as in vivo methods are concerned) are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/27728

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9119813 A	26-12-1991	AT 160821 T	15-12-1997
		AU 663053 B	28-09-1995
		AU 8206191 A	07-01-1992
		CA 2084987 A	12-12-1991
		DE 69128350 D	15-01-1998
		DE 69128350 T	26-03-1998
		DK 533838 T	23-02-1998
		EP 0533838 A	31-03-1993
		EP 0786469 A	30-07-1997
		ES 2110444 T	16-02-1998
		GR 3025594 T	31-03-1998
		IL 98456 A	18-03-1997
		IL 112141 A	05-04-1998
		JP 8252100 A	01-10-1996
		JP 2763958 B	11-06-1998
		KR 9702255 B	26-02-1997
		US 5459015 A	17-10-1995
		US 5707796 A	13-01-1998
		US 5472841 A	05-12-1995
		US 5543293 A	06-08-1996
		US 5849479 A	15-12-1998
		US 5683867 A	04-11-1997
		US 5503978 A	02-04-1996
		US 5527894 A	18-06-1996
		US 5650275 A	22-07-1997
		US 5637459 A	10-06-1997
		US 5648214 A	15-07-1997
		US 5723289 A	03-03-1998
		US 5629155 A	13-05-1997
		US 5756287 A	26-05-1998
		US 5641629 A	24-06-1997
		US 5639868 A	17-06-1997
		US 5622828 A	22-04-1997
		US 5705337 A	06-01-1998
		US 5637461 A	10-06-1997
		US 5696249 A	09-12-1997
		US 5670637 A	23-09-1997
		US 5660985 A	26-08-1997
		US 5750342 A	12-05-1998
		US 5688935 A	18-11-1997
		US 5763566 A	09-06-1998
		US 5712375 A	27-01-1998
		US 5789157 A	04-08-1998
		US 5864026 A	26-01-1999
		US 5637682 A	10-06-1997
		US 5734034 A	31-03-1998
		US 5595877 A	21-01-1997
		US 5587468 A	24-12-1996
		US 5786462 A	28-07-1998

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.